

1. NAME OF THE MEDICINAL PRODUCT

VALGEN-AM 160MG + 5MG FILM-COATED TABLETS

VALGEN-AM 160MG +10MG FILM-COATED TABLETS

2. NAME AND STRENGTH OF ACTIVE SUBSTANCES

VALGEN-AM 160MG + 5MG FILM-COATED TABLETS: Each tablet contains Valsartan 160 mg and Amlodipine besylate 6.935 mg equivalent to Amlodipine 5 mg

VALGEN-AM 160MG + 10MG FILM-COATED TABLETS: Each tablet contains Valsartan 160 mg and Amlodipine besylate 13.870 mg equivalent to Amlodipine 10 mg

3. PRODUCT DESCRIPTION

VALGEN-AM 160MG + 5MG FILM-COATED TABLETS

Dark Yellow, oval-shaped, biconvex, film-coated tablets debossed with 'L298' on one side and plain on other side.

VALGEN-AM 160MG + 10MG FILM-COATED TABLETS

Light Yellow, oval-shaped, biconvex, film-coated tablets debossed with 'L300' on one side and plain on other side.

4. DOSAGE FORM

Film-coated Tablets

5. PHARMACODYNAMIC PROPERTIES

Pharmacotherapeutic group: Angiotensin-II antagonists, plain (valsartan) combinations with dihydropyridine derivatives (amlodipine), ATC code: C09DB01.

Pharmacodynamics

VALGEN-AM combines two antihypertensive compounds with complementary mechanisms to control blood pressure in patients with essential hypertension: amlodipine belongs to the calcium antagonist class and valsartan to the angiotensin II (Ang II) antagonist class of medicines. The combination of these ingredients has an additive antihypertensive effect, reducing blood pressure to a greater degree than either component alone.

Amlodipine

The amlodipine component of VALGEN-AM inhibits the transmembrane entry of calcium ions into cardiac and vascular smooth muscle. The mechanism of the antihypertensive action of amlodipine is due to a direct relaxant effect on vascular smooth muscle, causing reductions in peripheral vascular resistance and blood pressure. Experimental data suggest that amlodipine binds to both dihydropyridine and non-dihydropyridine binding sites. The contractile processes of cardiac muscle and vascular smooth muscle are dependent upon the movement of extracellular calcium ions into these cells through specific ion channels.

Following administration of therapeutic doses to patients with hypertension, amlodipine produces vasodilatation resulting in a reduction of supine and standing blood pressures. These decreases in blood pressure are not accompanied by a significant change in heart rate or plasma catecholamine levels with chronic dosing.

Plasma concentrations correlate with effect in both young and elderly patients.

In hypertensive patients with normal renal function, therapeutic doses of amlodipine resulted in a decrease in renal vascular resistance and an increase in glomerular filtration rate and effective renal plasma flow without change in filtration fraction or proteinuria.

As with other calcium channel blockers, haemodynamic measurements of cardiac function at rest and during exercise (or pacing) in patients with normal ventricular function treated with amlodipine have generally demonstrated a small increase in cardiac index without significant influence on dP/dt or on left ventricular end diastolic pressure or volume. In haemodynamic studies, amlodipine has not been associated with a negative inotropic effect when administered in the therapeutic dose range to intact animals and humans, even when co-administered with beta blockers to humans.

Amlodipine does not change sinoatrial nodal function or atrioventricular conduction in intact animals or humans. In clinical studies in which amlodipine was administered in combination with beta-blockers to patients with either hypertension or angina, no adverse experiences on electrocardiographic parameters were observed.

Amlodipine has demonstrated beneficial clinical effects in patients with chronic stable angina, vasospastic angina and angiographically documented coronary artery disease.

Valsartan

Valsartan is an orally active, potent, and specific angiotensin II receptor antagonist. It acts selectively on the AT₁ receptor subtype, which is responsible for the known actions of angiotensin II. The increased plasma levels of angiotensin II following AT₁ receptor blockade with valsartan may stimulate the unblocked AT₂ receptor, which appears to counterbalance the effect of the AT₁ receptor. Valsartan does not exhibit any partial agonist activity at the AT₁ receptor and has much (about 20,000 fold) greater affinity for the AT₁ receptor than for the AT₂ receptor.

Valsartan does not inhibit ACE, also known as kininase II, which converts angiotensin I to angiotensin II and degrades bradykinin. Since there is no effect on ACE and no potentiation of bradykinin or substance P, angiotensin II antagonists are unlikely to be associated with cough. In clinical trials where valsartan was compared with an ACE inhibitor, the incidence of dry cough was significantly ($P < 0.05$) lower in patients treated with valsartan than in those treated with an ACE inhibitor (2.6% versus 7.9% respectively). In a clinical trial of patients with a history of dry cough during ACE inhibitor therapy, 19.5% of trial subjects receiving valsartan and 19.0% of those receiving a thiazide diuretic experienced cough compared to 68.5% of those treated with an ACE inhibitor ($P < 0.05$). Valsartan does not bind to or block other hormone receptors or ion channels known to be important in cardiovascular regulation.

Administration of Valsartan to patients with hypertension results in reduction of blood pressure without affecting pulse rate.

In most patients, after administration of a single oral dose, onset of antihypertensive activity occurs within 2 hours, and the peak reduction of blood pressure is achieved within 4-6 hours. The antihypertensive effect persists over 24 hours after administration. During repeated administration, the maximum reduction in blood pressure with any dose is generally attained within 2-4 weeks and is sustained during long-term therapy. Abrupt withdrawal of Valsartan has not been associated with rebound hypertension or other adverse clinical events.

In patients with chronic heart failure (NYHA class II-IV), valsartan has been demonstrated to significantly reduce hospitalizations in patients with chronic heart failure (NYHA class II-IV). The benefits were greatest in patients not receiving either an ACE inhibitor or a beta blocker. In post-MI patients, valsartan has also been shown to reduce cardiovascular mortality in clinically stable patients with left ventricular failure or left ventricular dysfunction following myocardial infarction.

6. PHARMACOKINETIC PROPERTIES

Linearity

Valsartan and amlodipine exhibit linear pharmacokinetics.

Amlodipine

Absorption

After oral administration of therapeutic doses of amlodipine alone, peak plasma concentrations of amlodipine are reached in 6 to 12 hours. Absolute bioavailability has been calculated as between 64% and 80%. Amlodipine bioavailability is unaffected by food ingestion.

Distribution

Volume of distribution is approximately 21 L/kg. *In vitro* studies with amlodipine have shown that approximately 97.5% of circulating drug is bound to plasma proteins. Amlodipine crosses the placenta and is excreted into breast milk.

Biotransformation

Amlodipine is extensively (approximately 90%) metabolized in the liver to inactive metabolites.

Elimination

Amlodipine elimination from plasma is biphasic with a terminal elimination half-life of approximately 30 to 50 hours. Steady-state plasma levels are reached after continuous administration for 7–8 days. Ten per cent of original amlodipine and 60% of amlodipine metabolites are excreted in urine.

Valsartan

Absorption

Following oral administration of valsartan alone, peak plasma concentrations of valsartan are reached in 2-4 hours. Mean absolute bioavailability is 23%. Food decreases the exposure (as measured by AUC) to valsartan by about 40% and peak plasma concentration (C_{max}) by about 50%, although from about 8 h post dosing plasma valsartan concentrations are similar for the fed and fasted group. This reduction in AUC, however, is not accompanied by a clinically significant reduction in therapeutic effect, and valsartan can therefore be given either with or without food.

Distribution

The steady-state volume of distribution of valsartan after intravenous administration is about 17 liters indicating that valsartan is not distributed into tissues extensively. Valsartan is highly bound to serum proteins (94-97%), mainly serum albumin.

Biotransformation

Valsartan is not transformed to a high extent as only about 20% of dose is recovered as metabolites. A hydroxy metabolite has been identified in plasma at low concentrations (less than 10 % of the valsartan AUC). This metabolite is pharmacologically inactive.

Elimination

Valsartan shows multiexponential decay kinetics ($1/2\alpha < 1\text{h}$ and $t_{1/2\beta}$ about 9 h). Valsartan is primarily eliminated unchanged in faeces (about 83% of dose) and urine (about 13% of dose) mainly as unchanged drug. Following intravenous administration, plasma clearance of valsartan is about 2 L/h and its renal clearance is 0.62 L/h (about 30% of total clearance). The half-life of valsartan is 6 hours.

Valsartan/Amlodipine

Following oral administration of VALGEN-AM peak plasma concentrations of valsartan and amlodipine are reached in 3 and 6-8 hours, respectively. The rate and extent of absorption of VALGEN-AM are equivalent to the bioavailability of valsartan and amlodipine when administered as individual tablets.

Special populations

Geriatric patients

The time to reach peak plasma concentrations of amlodipine is similar in elderly and younger subjects. Amlodipine clearance tends to be decreased with resulting increases in AUC and elimination half life in elderly patients.

Systemic exposure to valsartan is slightly elevated in the elderly as compared to the young, but this has not been shown to have any clinical significance.

Renal impairment

The pharmacokinetics of amlodipine is not significantly influenced by renal impairment. There is no apparent correlation between renal function (measured by creatinine clearance) and exposure (measured by AUC) to valsartan in patients with different degrees of renal impairment. Patients with mild to moderate renal impairment may therefore receive the usual initial dose.

Hepatic impairment

Patients with hepatic insufficiency have decreased clearance of amlodipine with resulting increase in AUC of approximately 40-60%. In a small number of patients with mild to moderate hepatic impairment given single doses of 5 mg, amlodipine half-life has been prolonged. Worsening of liver function test values may occur.

About 70% of the absorbed valsartan dose is excreted in the bile, mainly as unchanged compound. The AUC with valsartan has been observed to approximately double in patients with mild or moderate hepatic impairment including patients with biliary obstructive disorders. There are no data available on the use of valsartan in patients with severe hepatic dysfunction.

Care should be exercised in patients with liver disease.

7. INDICATIONS

Treatment of essential hypertension.

VALGEN-AM is indicated in patients whose blood pressure is not adequately controlled by monotherapy.

VALGEN-AM is indicated for the initial treatment of hypertension. The choice of VALGEN-AM for initial treatment should be based on an assessment of the potential benefits and risks.

8. DOSAGE REGIMEN AND ADMINISTRATION

DOSAGE REGIMEN

General target population

A patient whose blood pressure is not adequately controlled on monotherapy may be switched to combination therapy with VALGEN-AM. The recommended dose is one tablet per day. When clinically appropriate direct change from monotherapy to the fixed-dose combination may be considered.

For convenience, patients receiving valsartan and amlodipine from separate tablets may be switched to VALGEN-AM containing the same component doses.

For initial therapy the usual starting dose is Amlodipine and Valsartan tablets 5mg/80 mg once daily. The dosage can be increased after 1 to 2 weeks of therapy to a maximum of 10 mg / 320 mg per day as needed to control blood pressure. VALGEN-AM is not recommended as initial therapy in patients with intravascular volume depletion (Note: VALGEN-AM 5mg+80mg is not available in Malaysia).

The maximum dose is 10 mg / 320 mg per day.

Both amlodipine and valsartan monotherapy can be taken with or without food. It is recommended to take VALGEN-AM with some water.

The products are not able to deliver all approved dose regimen for the combination of amlodipine and valsartan tablet. Other approved dosage forms and strengths of the combination of amlodipine and valsartan tablet should be used in such cases.

Special populations

Geriatric patients (aged 65 years or above)

Since both components of the combination are equally well tolerated when used at similar doses in elderly (aged 65 years or above) or younger patients, no dose adjustment of the starting dose is required. Starting with the lowest available dose of amlodipine should be considered. The lowest strength of VALGEN-AM contains 5 mg of amlodipine.

Pediatric patients (below 18 years)

VALGEN-AM is not recommended for use in patients aged below 18 years due to a lack of data on safety and efficacy.

Renal impairment

No dosage adjustment is required for patients with mild to moderate renal impairment. Caution is required if severe renal impairment occur.

Hepatic impairment

Liver function should be monitored in patients with mild to moderate hepatic impairment. The daily dose of VALGEN-AM should not exceed 5/80mg in patients with mild to moderate hepatic impairment without cholestasis. VALGEN-AM is contraindicated in severe hepatic impairment. Starting with the lowest available dose of amlodipine should be considered. The lowest strength of VALGEN-AM contains 5 mg of amlodipine.

9. METHOD OF ADMINISTRATION

For oral use.

10. CONTRAINDICATIONS

- Known hypersensitivity to the amlodipine, valsartan or to any of the excipients.
- Pregnancy
- Severe hepatic impairment; biliary cirrhosis and cholestasis.
- Concomitant use of angiotensin receptor antagonists (ARBs) - including valsartan – or of angiotensin-converting enzyme inhibitors (ACEIs) with aliskiren in patients with Type 2 diabetes.

11. WARNING AND PRECAUTIONS

Patients with sodium- and/or volume depletion

Excessive hypotension was seen in 0.4% of patients with uncomplicated hypertension treated with VALGEN-AM in placebo-controlled studies. In patients with an activated renin- angiotensin system (such as volume- and/or salt-depleted patients receiving high doses of diuretics) who are receiving angiotensin receptor blockers, symptomatic hypotension may occur. Correction of this condition prior to administration of VALGEN-AM or close medical supervision at the start of treatment is recommended.

If hypotension occurs with VALGEN-AM, the patient should be placed in the supine position and, if necessary, given an I.V. infusion of normal saline. Treatment can be continued once blood pressure has been stabilized.

Hyperkalemia

Concomitant use with potassium supplements, potassium sparing diuretics, salt substitutes containing potassium, or other drugs that may increase potassium levels (heparin, etc.) should be used with caution and with frequent monitoring of potassium.

Patients with renal artery stenosis

VALGEN-AM should be used with caution to treat hypertension in patients with unilateral or bilateral renal artery stenosis, stenosis to a solitary kidney since blood urea and serum creatinine may increase in such patients.

Patients with renal impairment

No data is available for severe cases (creatinine clearance < 10 mL/min.) and caution is therefore advised. No dosage adjustment of VALGEN-AM is required for patients with mild to moderate renal impairment.

The use of ARBs - including valsartan - or of ACEIs with aliskiren should be avoided in patients with severe renal impairment (GFR < 30 mL/min).

Patients with kidney transplantation

To date there is no experience of the safe use of VALGEN-AM in patients who have had recent kidney transplantation.

Patients with hepatic impairment

Valsartan is mostly eliminated unchanged via the bile whereas amlodipine is extensively metabolized by the liver. In patient with mild to moderate hepatic impairment without cholestasis, VALGEN-AM should be used with caution and careful monitoring of liver function tests should be performed. The daily dose of VALGEN-AM should not exceed 5/80mg. Patients with severe hepatic impairment, biliary cirrhosis or cholestasis should not take VALGEN-AM.

Angioedema

Angioedema, including swelling of the larynx and glottis, causing airway obstruction and/or swelling of the face, lips, pharynx, and/or tongue has been reported in patients treated with valsartan; some of these patients previously experienced angioedema with other drugs including ACE inhibitors. VALGEN-AM should be immediately discontinued in patients who develop angioedema, and VALGEN-AM should not be re-administered.

Patients with heart failure/post-myocardial infarction

In general, calcium channel blockers including amlodipine should be used with caution in patients with serious congestive heart failure (New York Heart Association (NYHA) functional class III-IV).

In patients whose renal function may depend on the activity of the renin-angiotensin- aldosterone system (e.g. patients with severe congestive heart failure), treatment with angiotensin converting enzyme inhibitors or angiotensin receptor antagonists has been associated with oliguria and/or progressive azotemia, and in rare cases with acute renal failure and/or death. Evaluation of patients with heart failure or post-myocardial infarction should always include assessment of renal function.

Patients with acute myocardial infarction

Worsening angina pectoris and acute myocardial infarction can develop after starting or increasing the dose of amlodipine, particularly in patients with severe obstructive coronary artery disease.

Patients with aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy

As with all other vasodilators, special caution is required when using amlodipine in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy.

Dual Blockade of the Renin-Angiotensin-Aldosterone System (RAAS)

Caution is required while co-administering ARBs, including valsartan, with other agents blocking the RAAS such as ACEIs or aliskiren. (See section INTERACTION, Subsection Dual Blockade of the RAAS)

12. INTERACTIONS WITH OTHERMEDICAMENTS

Amlodipine

Simvastatin: Co-administration of multiple doses of 10 mg of amlodipine with 80 mg simvastatin resulted in a 77% increase in exposure to simvastatin compared to simvastatin alone. It is recommended to limit the dose of simvastatin to 20 mg daily in patients on amlodipine.

CYP3A4 Inhibitors: Co-administration of a 180 mg daily dose of diltiazem with 5 mg amlodipine in elderly hypertensive patients resulted in a 1.6 fold increase in amlodipine systemic exposure. However, strong inhibitors of CYP3A4 (e.g., ketoconazole, itraconazole, ritonavir) may increase the plasma concentrations of amlodipine to a greater extent than diltiazem. Caution should therefore be exercised when co-administering amlodipine with CYP3A4 inhibitors.

Grapefruit Juice: The exposure of amlodipine may be increased when co-administered with grapefruit juice due to CYP3A4 inhibition. However, co-administration of 240 mL of grapefruit juice with a single oral dose of amlodipine 10 mg in 20 healthy volunteers had no significant effect on the pharmacokinetics of amlodipine.

CYP3A4 Inducers: No information is available on the quantitative effects of CYP3A4 inducers on amlodipine. Patients should be monitored for adequate clinical effect when amlodipine is co-administered with CYP3A4 inducers (e.g. rifampicin, hypericum perforatum).

In monotherapy, amlodipine has been safely administered with thiazide diuretics, beta- blockers, angiotensin-converting enzyme inhibitors, long-acting nitrates, sublingual nitroglycerin, digoxin, warfarin, atorvastatin, sildenafil, maalox (Aluminium hydroxide gel, Magnesium hydroxide and Simeticone), cimetidine, non-steroidal anti-inflammatory drugs, antibiotics, and oral hypoglycemic drugs.

Valsartan

Dual blockade of the Renin-Angiotensin-Aldosterone System (RAAS) with ARBs, ACEIs or aliskiren:

The concomitant use of ARBs, including valsartan, with other agents acting on the RAAS is associated with an increased incidence of hypotension, hyperkalaemia, and changes in renal function compared to monotherapy.

It is recommended to monitor blood pressure, renal function and electrolytes in patients on VALGEN-AM and other agents that affect the RAAS.

The concomitant use of ARBs – including valsartan - or of ACEIs with aliskiren, should be avoided in patients with severe renal impairment (GFR < 30 mL/min).

The concomitant use of ARBs including valsartan, or ACEIs, with aliskiren is contraindicated in patients with Type 2 diabetes mellitus.

Potassium: Concomitant use with potassium supplements, potassium-sparing diuretics, salt substitutes containing potassium, or other drugs that may increase potassium levels (heparin, etc.) requires caution and frequent monitoring of potassium levels.

Non-Steroidal Anti-Inflammatory Agents (NSAIDs) including Selective Cyclooxygenase-2 Inhibitors (COX-2 Inhibitors): When angiotensin II antagonists are administered simultaneously with NSAIDs, attenuation of the antihypertensive effect may occur. Furthermore, in patients who are elderly, volume-depleted (including those on diuretic therapy), or have compromised renal function, concomitant use of angiotensin II antagonists and NSAIDs may lead to an increased risk of worsening of renal function. Therefore, monitoring of renal function is recommended when initiating or modifying the treatment in patients on valsartan who are taking NSAIDs concomitantly.

Lithium: Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with ACE inhibitors or angiotensin II receptor antagonists including VALGEN-AM. Therefore, careful monitoring of serum lithium levels is recommended during concomitant use. If a diuretic is also used, the risk of lithium toxicity may presumably be increased further with VALGEN-AM.

Transporters: The results from an *in vitro* study with human liver tissue indicate that valsartan is a substrate of the hepatic uptake transporter OATP1B1 and the hepatic efflux transporter MRP2. Co-administration of inhibitors of the uptake transporter (e.g. rifampin, cyclosporin) or efflux transporter (e.g. ritonavir) may increase the systemic exposure to valsartan.

In monotherapy with valsartan, no drug interactions of clinical significance have been found with the following drugs: cimetidine, warfarin, furosemide, digoxin, atenolol, indomethacin, hydrochlorothiazide, amlodipine, glibenclamide.

13. PREGNANCY AND LACTATION

Pregnancy

Risk Summary

As for any drug that acts directly on the renin-angiotensin-aldosterone system (RAAS), VALGEN-AM must not be used during pregnancy. Due to the mechanism of action of angiotensin II antagonists, a risk to the foetus cannot be excluded.

Administration of angiotensin converting enzyme (ACE) inhibitors (a specific class of drugs acting on the renin-angiotensin-aldosterone system (RAAS)) to pregnant women during the second and third trimesters has been reported to cause injury and death to the developing foetus. In addition, in retrospective data, first trimester use of ACE inhibitors has been associated with a potential risk of birth defects. There have been reports of spontaneous abortion, oligohydramnios and new born renal dysfunction when pregnant women have inadvertently taken valsartan.

There are no adequate clinical data with amlodipine in pregnant women. Animal studies with amlodipine have shown reproductive toxicity at dose 8 times the maximum recommended dose of 10 mg. The potential risk to humans is unknown.

If pregnancy is detected during therapy, VALGEN-AM must be discontinued as soon as possible.

Clinical considerations

Disease-associated maternal and/or embryo/fetal risk

Hypertension in pregnancy increases the maternal risk for pre-eclampsia, gestational diabetes, premature delivery, and delivery complications (e.g., need for cesarean section, and post-partum hemorrhage). Hypertension increases the fetal risk for intrauterine growth restriction and intrauterine death.

Fetal/Neonatal Risk

Oligohydramnios in pregnant women who use drugs affecting the renin-angiotensin system in the second and third trimesters of pregnancy can result in the following: reduced fetal renal function leading to anuria and renal failure, fetal lung hypoplasia, skeletal deformations, including skull hypoplasia, hypotension and death.

In case of accidental exposure to ARB therapy, appropriate fetal monitoring should be considered.

Infants whose mothers have taken ARB therapy in the first trimester should be closely observed for hypotension.

Lactation

It is not known whether valsartan is excreted in human milk. It is reported that amlodipine is excreted in human milk. The proportion of the maternal dose received by the infant has been estimated with an interquartile range of 3–7%, with a maximum of 15%. The effect of amlodipine on infants is unknown. Valsartan was excreted in the milk of lactating rats. It is therefore not advisable for women who are breast-feeding to use VALGEN-AM.

Females and males of reproductive potential

As for any drug that acts directly on the renin-angiotensin-aldosterone system (RAAS), VALGEN-AM must not be used in women planning to become pregnant. Healthcare professionals prescribing any agents acting on the renin-angiotensin-aldosterone system (RAAS) should counsel women of childbearing potential about the potential risk of these agents during pregnancy.

Infertility

There is no information on the effects of amlodipine or valsartan on human fertility. Studies in rats did not show any effects of amlodipine or valsartan on fertility.

14. ADVERSE EFFECTS / UNDESIRABLE EFFECTS

The safety of VALGEN-AM has been evaluated in five controlled clinical studies with patients of

whom received valsartan in combination with amlodipine.

Adverse drug reactions or adverse experiences (Table 1 and Table 2) are ranked under heading of frequency, the most frequent first, using the following convention: very common ($\geq 1/10$); common ($\geq 1/100$, $< 1/10$); uncommon ($\geq 1/1,000$, $< 1/100$); rare ($\geq 1/10,000$, $< 1/1,000$) very rare ($< 1/10,000$), including isolated reports. Within each frequency grouping, adverse reactions are ranked in order of decreasing seriousness.

Table 1: Adverse drug reactions with VALGEN-AM

Infections and infestations	
Common:	Nasopharyngitis, influenza
Immune system disorders	
Rare:	Hypersensitivity
Eye disorders	
Rare	Visual disturbance
Psychiatric disorders	
Rare:	Anxiety
Nervous system disorders	
Common:	Headache
Uncommon:	Dizziness, somnolence, dizziness postural, paresthesia
Ear and labyrinth disorders	
Uncommon:	Vertigo
Rare:	Tinnitus
Cardiac disorders	
Uncommon:	Tachycardia, palpitations
Rare:	Syncope
Vascular Disorders	
Uncommon:	Orthostatic hypotension
Rare:	Hypotension
Respiratory, thoracic and mediastinal disorders	
Uncommon:	Cough, pharyngolaryngeal pain
Gastrointestinal disorders	
Uncommon:	Diarrhea, nausea, abdominal pain, constipation, dry mouth
Skin and subcutaneous tissue disorders	
Uncommon:	Rash, erythema
Rare:	Hyperhidrosis, exanthema, pruritus
Musculoskeletal and connective tissue disorders	
Uncommon:	Joint swelling, back pain, arthralgia
Rare:	Muscle spasm, sensation of heaviness
Renal and urinary disorders	
Rare:	Pollakiuria, polyuria

Reproductive system and breast disorders

Rare: Erectile dysfunction

General disorders and administration site conditions

Common: Oedema, pitting oedema, facial oedema, oedema peripheral, fatigue, flushing, asthenia, hot flush

Additional information on the combination

In double-blind, active- or placebo-controlled completed clinical trials, the incidence of peripheral edema was statistically lower in patients treated with the combination (5.8%) than in patients treated with amlodipine monotherapy (9%).

Laboratory evaluation

Very few hypertensive patients treated with valsartan/amlodipine showed notable changes in laboratory test results from baseline. There was a slightly higher incidence of notably increased blood urea nitrogen in the amlodipine/valsartan (5.5 %) and valsartan monotherapy (5.5%) groups as compared to the placebo group (4.5%).

Additional information on individual components

Adverse reactions previously reported with one of the individual components may occur with VALGEN-AM even if not observed in clinical trials.

Amlodipine

Other additional adverse experiences reported in clinical trials with amlodipine monotherapy, irrespective of their causal association with the study drug, are presented in Table 2: Because amlodipine clinical trials were conducted under widely varying conditions, adverse experience rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Table 2: Adverse experiences with amlodipine monotherapy

Eye disorders	
Uncommon	Diplopia
Blood and lymphatic system disorders	
Very rare	Thrombocytopenia, leucocytopenia
Immune system disorders	
Very rare	Allergic reactions
Metabolism and nutrition disorders	
Very rare	Hyperglycemia
Psychiatric disorders	
Uncommon	Insomnia, mood changes
Nervous system disorders	

Uncommon	Tremor, hypoesthesia, dysgeusia
Very rare	Peripheral neuropathy, hypertonia
Cardiac disorders	
Very rare	Arrhythmia, bradycardia, atrial fibrillation, ventricular tachycardia, myocardial infarction
Vascular disorders	
Very rare	Vasculitis
Respiratory, thoracic and mediastinal disorders	
Uncommon	Dyspnea, rhinitis
Gastrointestinal disorders	
Uncommon	Vomiting, dyspepsia
Very rare	Pancreatitis, gastritis, gingival hyperplasia
Hepatobiliary disorders	
Very rare	Hepatitis, jaundice
Skin and subcutaneous tissue disorders	
Uncommon	Alopecia, purpura, skin discoloration, photosensitivity
Very rare	Angioedema, urticaria, erythema multiforme, Steven Johnson syndrome
Musculoskeletal and connective tissue disorders	
Uncommon	Myalgia
Renal and urinary disorders	
Uncommon	Micturition disorder, nocturia
Reproductive system and breast disorders	
Uncommon	Gynecomastia
General disorders and administration site conditions	
Uncommon	Pain, malaise, chest pain
Investigations	
Uncommon	Weight decreased, weight increased
Very rare	Hepatic enzyme increased(mostly consistent with cholestasis)

Valsartan

Other ADRs reported from clinical studies, post-marketing experience and laboratory findings in hypertension indication are presented in Table 3 according to system organ class.

For all the ADRs reported from post-marketing experience and laboratory findings, it is not possible to apply any ADR frequency and therefore they are mentioned with a "not known" frequency.

Table 3: Adverse drug reactions with valsartan monotherapy

Blood and lymphatic system disorders	
Not known	Hemoglobin decreased, Hematocrit decreased, Neutropenia, Thrombocytopenia
Immune system disorders	
Not known	Hypersensitivity including serum sickness
Metabolism and nutrition disorders	
Not known	Blood potassium increased
Vascular disorders	
Not known	Vasculitis
Hepato-biliary disorders	
Not known	Liver function test abnormal including blood bilirubin
Skin and subcutaneous tissue disorders	
Not known	Angioedema, dermatitis bullous
Musculoskeletal and connective tissue disorders	
Not known	Myalgia
Renal and urinary disorders	
Not known	Renal failure and impairment, Blood creatinine increased

The following events have also been observed during clinical trials in hypertensive patients irrespective of their causal association with the study drug: Insomnia, libido decrease, pharyngitis, rhinitis, sinusitis, upper respiratory tract infection, viral infections.

15. OVERDOSE AND TREATMENT

Symptom

There is no experience of overdose with VALGEN-AM yet. The major symptom of overdose with valsartan is possibly pronounced hypotension with dizziness. Overdose with amlodipine may result in excessive peripheral vasodilatation and possibly reflex tachycardia. Marked and potentially prolonged systemic hypotension up to and including shock with fatal outcome have been reported.

Treatment

Clinically significant hypotension due to amlodipine overdose calls for active cardiovascular support including frequent monitoring of cardiac and respiratory function, elevation of extremities, and attention to circulating fluid volume and urine output.

A vasoconstrictor may be helpful in restoring vascular tone and blood pressure, provided that there is no contraindication to its use. If the ingestion is recent, induction of vomiting or gastric lavage may be considered.

Administration of activated charcoal to healthy volunteers immediately or up to two hours after ingestion of amlodipine has been shown to significantly decrease amlodipine absorption.

Intravenous calcium gluconate may be beneficial in reversing the effects of calcium channel blockade.

Both valsartan and amlodipine are unlikely to be removed by haemodialysis.

16. EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Patient taking VALGEN-AM and driving vehicles or using machines should take into account that dizziness or weariness may occasionally occur. Amlodipine can have a mild or moderate influence on the ability to drive and use machines. If patients taking amlodipine suffer from dizziness, headache, fatigue or nausea the ability to react may be impaired.

17. STORAGE CONDITION

Store below 30°C.

Protect from moisture. Store in the original package.

18. PACKAGING AVAILABLE

Alu/ Alu blisters of 2 x 8's, 4 x 8's, 7 x 8's, and 12 x 8's in blister form packed in the outer carton. Not all pack sizes may be marketed.

19. NAME AND ADDRESS OF MANUFACTURER / PRODUCT REGISTRATION

Manufactured by:

Alembic Pharmaceuticals Limited, (Formulation Division)
Village Panelav, P.O. Tajpura, Near Baska,
Taluka Halol, Panchmahal, Gujarat-389350 India.

Product Registration Holder:

GENPHARMA SDN. BHD.
Lot 5016, Jalan Teratai,
5 ½ Mile Off Jalan Meru, 41050 Klang,
Selangor, Malaysia

20. DATE OF REVISION

22/03/24

(Font size:11, as measure in Times New Roman)