

Acknowledgment



Draft

Glucocorticoid Preparation

Shincort

Injection 10mg



Each mL contains:
Triamcinolone Acetonide 10mg
Benzyl Alcohol (as preservative) 0.02mL
Propylene Glycol (as preservative) 0.2mL

5mL x 10Vials

Each mL contains: Triamcinolone Acetonide 10mg
Preservatives: Benzyl Alcohol 0.02mL, Propylene Glycol 0.2mL
Indications: Rheumatoid arthritis, synovitis, allergic rhinitis, allergic asthma, allergic dermatitis, pemphigus, allergic and inflammatory ophthalmic diseases.
Dosage and Administration: Recommended dosage: For adults and children over 12 years of age, the suggested initial dose is 60 mg, injected deeply into the gluteal muscle. Dosage is usually adjusted within the range of 40 to 80mg. See insert for more information. To be administered by a physician.
Route of Administration: Intramuscular, intra-articular and intrabursal administration.
Storage conditions: Store at temperature not exceeding 30°C. Avoid freezing and protect from light.

SHI10 MYS-2167069-013

Keep medicine out of reach of children.
Jauhkan daripada kanak-kanak.

Controlled Medicine / Ubat Terkawal

Importer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199201001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info Line: 1 800 88 3679

Warning: As this preparation contains benzyl alcohol, its use should be avoided in children under two years of age. Not to be used in neonates.

R.O.C. Reg. No.: 11476 Batch No.: SHI10
DCA Reg. No.: MAL 19890671AZ
MYN R1708A5618 Mfg. Date:
SIN 6709P Exp. Date:
BRU11120833P

Manufactured by:
YUNG SHIN PHARMACEUTICAL IND. CO., LTD.

Taichung Youth Factory
No. 27, Kung Chiu Rd., Tachia, Taichung,
Taiwan, R.O.C.

A Subsidiary of **Y.S.P.SAH®**

150%

Glucocorticoid Preparation

Shincort

Injection 10mg



Each mL contains:
Triamcinolone Acetonide 10mg
Benzyl Alcohol (as preservative) 0.02mL
Propylene Glycol (as preservative) 0.2mL

5mL x 10Vials

Each mL contains: Triamcinolone Acetonide 10mg
Preservatives: Benzyl Alcohol 0.02mL, Propylene Glycol 0.2mL
Indications: Rheumatoid arthritis, synovitis, allergic rhinitis, allergic asthma, allergic dermatitis, pemphigus, allergic and inflammatory ophthalmic diseases.
Dosage and Administration: Recommended dosage: For adults and children over 12 years of age, the suggested initial dose is 60 mg, injected deeply into the gluteal muscle. Dosage is usually adjusted within the range of 40 to 80mg. See insert for more information. To be administered by a physician.
Route of Administration: Intramuscular, intra-articular and intrabursal administration.
Storage conditions: Store at temperature not exceeding 30°C. Avoid freezing and protect from light.

SHI10 MYS-2167069-013

Keep medicine out of reach of children.
Jauhkan daripada kanak-kanak.

Controlled Medicine / Ubat Terkawal

Importer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info Line: 1 800 88 3679

Warning: As this preparation contains benzyl alcohol, its use should be avoided in children under two years of age. Not to be used in neonates.

R.O.C. Reg. No.: 11476 Batch No.: SHI10
DCA Reg. No.: MAL 19890671AZ
MYN R1708A5618 Mfg. Date:
SIN 6709P Exp. Date:
BRU11120833P

Manufactured by:
YUNG SHIN PHARMACEUTICAL IND. CO., LTD.

Taichung Youth Factory
No. 27, Kung Chiu Rd., Tachia, Taichung,
Taiwan, R.O.C.

A Subsidiary of **Y.S.P.SAH®**