

YSP IBUPROFEN FILM COATED TABLET

Ibuprofen (400mg)

What is in this leaflet

1. What IBUPROFEN F. C. TABLET is used for.
2. How IBUPROFEN F. C. TABLET works.
3. Before you use IBUPROFEN F. C. TABLET
4. How to use IBUPROFEN F. C. TABLET
5. While you are using it
6. Side effects.
7. Storage and disposal of IBUPROFEN F. C.
8. Product description
9. Manufacturer and Product Registration Holder.
10. Date of revision

What IBUPROFEN F. C. TABLET is used for

IBUPROFEN F. C. TABLET relieves inflammation and pain caused by rheumatoid arthritis (systemic inflammatory disorder that affects joints) and osteoarthritis (degenerative joint disease). It can be used to relieve muscular and period pain too.

How IBUPROFEN F. C. TABLET works

Ibuprofen F.C. tablet contains a substance called Ibuprofen. Ibuprofen is a non-steroidal anti-inflammatory agent (NSAIDs) that has analgesic (pain-killing).

Before you use IBUPROFEN F. C. TABLET

- When you must not take it

Do not use Ibuprofen F.C. tablet: if you are allergic or hypersensitivity to Ibuprofen or any other ingredient listed at the end of this leaflet.

If you have experienced hypersensitivity reactions such as symptoms of of nasal polyps, angioedema (swelling around the eyes and lips and sometimes of the genitals, hands, and feet) and bronchospastic reactivity (constriction of the airway) to Aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs)

Do not use this medicine if any of the above condition applies to you. If you are not sure, get advice from your doctor or pharmacist

- Before you start to take it

Tell your doctor:

1. if you have or have had history of ulceration, bleeding and perforation of the stomach and intestinal.

2. if you are elderly
3. if you are smoking or drinking alcohol
4. if you have heart disease or stroke
5. if you have high blood pressure
6. if you have fluid retention or heart failure
7. Ibuprofen may cause severe skin reactions, e.g. exfoliative dermatitis, toxic epidermal necrolysis, Steven-Johnson Syndrome. Symptoms may include skin rash, blisters and peel off. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.
8. if you have bleeding disorder
9. if you have blood disorder or you are taking oral anticoagulant (blood thinning agents)
10. if you have infection
11. if you have liver or kidney disease
12. Please consult your doctor before start to use Ibuprofen F.C. tablet if you are pregnant, plan to become pregnant, breast-feeding or intend to breast-feed. Your doctor will have weighed the risks of you taking Ibuprofen F.C. tablet against the benefits they expect it will have for you.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines when taken with Ibuprofen F.C. tablet may interfere with each other. These include:

1. Hydantoins, medicines for fits
2. Sulfonamides, an antibiotic
3. Sulfonyleureas, for diabetes
4. anticoagulants, blood thinning agents
5. Aspirin, medicine for heart
6. other NSAIDs

How to use IBUPROFEN F. C. TABLET

- How much to use

IBUPROFEN F. C. TABLET should be taken according to prescription by the doctor.

- When to use it

IBUPROFEN F. C. TABLET should be taken according to the time advised by your doctor or pharmacist.

Do not exceed 3200mg total daily dose. If stomach and intestinal complaints occur, administer Ibuprofen with meals or milk.

Analgesic (pain relief) & antipyretic (to reduce fever): 600 - 1200mg daily, in divided doses.

For rheumatoid arthritis and osteoarthritis (joint disease): 1200 - 3200mg daily in divided doses.

Dysmenorrhea (period pain): 400mg every 4 hours when necessary for relief of pain.

A suggested dose for children:

20mg/kg body weight daily, in divided doses.

Not recommended for children under age of 1 year or for children weighing less than 7 kg. For short term use only.

- How long to use it

IBUPROFEN F. C. TABLET should be taken as your doctor advised you to do so.

- If you forget to use it

If you miss a dose of IBUPROFEN F. C. TABLET, take the missed dose as soon as you remember. However, if it's almost time for the next dose, skip the missed dose and continue the regular dosing schedule. Do not take a double dose to make up for a missed one.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have apply/taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking product name.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give this medication to anyone else, even if they have the same condition as you.

- Things to be careful of

Be careful driving or operating machinery until you know how IBUPROFEN F.C. TABLET affects you.

Inactive Ingredients-

Sodium starch glycolate, hydroxypropyl cellulose-L, talc, magnesium stearate, aerosil #200, hydroxypropyl methylcellulose, polyethylene glycol, dimethylpolysiloxane, purified water.

- MAL no.

MAL 19930602AZ

Side effects

Possible side effects include nausea, epigastric pain (abdominal pain), heartburn, and diarrhea. Other side effects that might appear such as dizziness, dyspepsia, headache, nervousness, pruritus (skin itching), tinnitus (ringing in the ears), depression, insomnia, blurred vision. Side effects are thrombocytopenia (low blood platelet count), agranulocytosis (low white blood cell count), hypersensitivity and liver and kidney function disorder.

If you encounter any other side effects not listed here, please consult your doctor immediately.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Manufacturer

Y.S.P Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7,
Section 13, Kawasan Perindustrian
Bandar Baru Bangi, 43000 Kajang,
Selangor, Malaysia.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision :

06/12/2025

Serial Number:

BPFK(R4/1)031215/00480

Storage and Disposal of IBUPROFEN F. C.- Storage

Store below 30°C, protect from light & moisture.

- Disposal

Dispose when you stop taking the tablets as instructed by your doctor or pharmacist or the tablets have passed their expiry date. Ask them what to do with any tablets that are left over.

Product description- What it looks like

A white color elliptical film coated tablet, both sides impressed with a score.

“”.

- Ingredients

Active Ingredient-

Ibuprofen