

CINNA[®] TABLET

Cinnarizine is a new type of vasodilator in the treatment of consequences of cerebral and coronary arteriosclerosis. It also has the properties and uses of the antihistamines, such as Meclizine, but it is mainly used for the symptomatic treatment of nausea and vertigo due to Meniere's disease and other labyrinthine disturbances.

Ingredient(s):

Each tablet contains:

Cinnarizine 25mg

Pharmacodynamics:

1. Cinnarizine increases and normalizes cerebral circulation and thereby increase the oxygen supply of the arteriosclerotic cerebral vessels. It prevents and eliminates vascular spasms of the labyrinth.
2. The improvement of cerebral circulation is also sustained by the fact that Cinnarizine increases the blood flow as well as the blood and oxygen supply to the heart muscle. The heart capacity is improved.
3. The mechanism of action of Cinnarizine may be explained by its direct inhibitory action on the tonus of the smooth muscles of certain arterioles which renders less sensitive to certain contractile substances such as angiotensin.

Pharmacokinetics:

It acts within about 30 minutes after oral administration, and its effect last for about 4 hours, with peak level achieved with 2 ~ 3 hours. About 90 % of the drug is excreted in the urine and stool after 72 hours of oral administration.

Indication(s):

For the symptomatic treatment of nausea and vertigo due to Meniere's disease and other labyrinthine disturbances; for the prevention of motion sickness; also used in the management of peripheral arterial disease.

Dosage and Administration:

The usual dose is 15 - 30mg three times daily.

Children aged 5 to 12 years may be given half the adult dose.

For motion sickness: 30mg may be taken 2 hours before the start of the journey and 15mg every 8 hours during the journey.

Mode of administration:

To be taken orally

Symptoms and Treatment for Overdosage and Antidote(s):

It is nontoxic substance, only doses may cause slight sleepiness, which disappears rapidly after reducing the dose.

Precaution(s) / Warning(s):

1. May cause drowsiness, patients affected in this way should not drive or operate machinery.
2. Avoid alcohol drink.
3. It should be used with reasonable caution in hypotensive patients (It has been found to decrease blood pressure significantly).
4. It is not advisable to administer in pregnant women.

Side Effect(s) / Adverse Reaction(s):

Only high doses may occasionally cause a slight sleepiness which disappears rapidly after reducing the dose. Other side effects are skin allergic reaction and fatigue.

Interaction with Other Medicaments:

May enhance sedative effects of CNS depressant. Atropine-like effects potentiated by other anticholinergic, tricyclic antidepressant and MAOIs.

Pregnancy and Lactation:

It is not advisable to administer in pregnant women.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description:

A white color round tablet one side is impressed with a score and the other side with "C".

**Packing(s):**

Plastic bottle of 100's, 200's, 250's, 500's, 1000's and 2000's (for export only).

Blister packing of 10's x 10 and 10's x 100.

The information contained in this leaflet is limited.
Further information can be obtained from other
sources including your doctor or pharmacist.

CINNA[®] TABLET

Bahan Aktif:

Setiap tablet mengandungi:

Cinnarizine 25mg

Indikasi:

Rawatan simptomatik bagi mual dan vertigo akibat penyakit Meniere dan gangguan labirin. Mencegah mabuk perjalanan. Juga untuk penyakit arteri perifer.

Dos:

Biasanya 15 - 30mg 3 kali sehari.

Kanak-kanak berumur 5 - 12 tahun separuh dos orang dewasa.

Mabuk perjalanan: 30mg diambil 2 jam sebelum perjalanan, kemudian 15mg setiap 8 jam sepanjang perjalanan.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679