


Item Code	: 2181093 - 09 draft 15 Aug 2025
PM Code	: Case - GFT 4's x 25 V1
Measurement(mm) ± 2mm	: 155 (L) x 105 (W) x 127 (H)
Colour of Artwork	: Pantone 2583 C Pantone 106 C K
Source of Artwork	: MAC 2 / CC

This design also applicable for pack size 4's x 150



**Ingredients:**  
 Each lozenge contains:  
 Chlorhexidine HCl ..... 5mg  
 Benzocaine ..... 2mg

**Flavoring Agent:**  
 Spearmint Oil ..... 0.025mg  
 Peppermint Oil ..... 0.025mg

**Indication(s):**  
 For the prophylaxis of stomatitis, gangrenous stomatitis, pharyngitis, laryngitis, tonsillitis, gingivitis, post-tonsillectomy and post tooth extraction secondary bacterial infections. For the temporary local relief of pain associated with dental conditions and sore throats.

**Dosage and Administrations:**  
 For prevention: 1 lozenge 4 - 5 times daily.  
 Place the lozenge on the tongue and let it dissolve slowly for longer bactericidal action.  
 To be dispensed on physician's prescription.

**Route of Administration:**  
 Oral.

This product contains chlorhexidine. Inform your healthcare provider if you have a known allergy to chlorhexidine.  
 Stop using this product and seek immediate medical assistance if you experience rash, itching, swelling, breathing difficulties, light-headedness or rapid heartbeat.

KEEP MEDICINE OUT OF REACH OF CHILDREN.  
 STORE AT TEMPERATURE BELOW 30°C.  
 PROTECT FROM LIGHT AND MOISTURE.

UBAT TERKAWAL  
 CONTROLLED MEDICINE

Manufacturer and Product Registration Holder:  
**Y.S.P. INDUSTRIES (M) SDN. BHD.**, (199001001034)  
 Lot 3, 5, 7, 12 & 14, Jalan P17, Section 13,  
 Kawasan Perindustrian Bandar Baru Bangi,  
 43000 Kajang, Selangor, Malaysia.  
 Ordering Line: 1 800 88 3027  
 Product Info: 1 800 88 3679

Distributor in Ethiopia:  
**West Pharma Trading PLC.**

Scan QR code to access package insert

**Bahan Aktif:**  
 Setiap lozeng mengandungi:  
 Chlorhexidine HCl ..... 5mg  
 Benzocaine ..... 2mg

**Perisa:**  
 Spearmint Oil ..... 0.025mg  
 Peppermint Oil ..... 0.025mg

**Indikasi:**  
 Untuk profilaksi bagi stomatitis, stomatitis bergangren, faringitis, laringitis, tonsilitis, gingivitis, jangkitan bakteria sekunder selepas tonsilektomi dan pencabutan gigi. Untuk pelegaan sementara sakit gigi dan sakit tekak.

**Dos:**  
 Untuk pencegahan: 1 lozeng, 4-5 kali sehari. Letakkan lozeng di atas lidah dan biarkan ia larut. Digunakan mengikut arahan doktor.

JAUHKAN DARIPADA KANAK-KANAK.  
 SIMPAN PADA SUHU BAWAH 30°C.  
 LINDUNGI DARIPADA CAHAYA DAN KELEMBAPAN

MAL19940289AZ  
 MYN R2601A1760  
 BRU18092416P

K1000 Trading  
 CAM R, 3220P-05

9 557892 100482

Batch No.:  
 Mfg. Date:  
 Exp. Date:

W  
 105mm  
 (± 2mm)

L  
 155mm  
 (± 2mm)

H  
 127mm  
 (± 2mm)