

DOMPER® TABLET

Domperidone (10mg)

What is in this leaflet

1. What DOMPER® TABLET is used for
2. How DOMPER® TABLET works
3. Before you use DOMPER® TABLET
4. How to use DOMPER® TABLET
5. While you are using it
6. Side effects
7. Storage and Disposal of DOMPER® TABLET
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What DOMPER® TABLET is used for

DOMPER® TABLET is indicated for the symptomatic relief of nausea and vomiting.

How DOMPER® TABLET works

DOMPER® TABLET is an antiemetic and a prokinetic medicine. It works by blocking the action of a chemical messenger in the brain which causes the feeling of nausea and vomiting, as well as increasing the movement or contractions of the stomach and intestines, allowing food to move more easily through the stomach.

Before you use DOMPER® TABLET

- When you must not use it

Do not take this medicine if:

- You are allergic to domperidone or any other ingredients of this medicine (listed in Product Description)
- You have moderate to severe liver impairment
- You have a tumour of the pituitary gland known as prolactinoma
- You have a pre-existing heart condition.
- You have gastrointestinal hemorrhage or mechanical obstruction.

- Before you start to use it

Please inform your healthcare professional regarding the following:

- You are pregnant or planning to become pregnant.
- Your baby is under 1 year of age.
- You are breast feeding or wish to breastfeed. (It is not advisable to take DOMPER® TABLET in

pregnancy because domperidone is found in the breast milk but at very low levels)

- You have or have ever had liver or kidney disease.
- You are not able to digest lactose which is a sugar found in milk and milk products.

Your doctor will advise you whether DOMPER® TABLET is suitable for you or if you need to adjust the dose.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

DOMPER® TABLET may interfere with some medicines. In particular, tell your healthcare professional if you are taking any of the following medicines listed below:

- Medicines for treating fungal infections, such as azole antifungals
- Antibiotic such as clarithromycin, levofloxacin, and erythromycin.
- Amiodarone, disopyramide, quinidine used to treat irregular heart rate.
- Medicines for treating mental disorder such as haloperidol, pimozide
- Antidepressants such as citalopram and escitalopram
- Cisapride used to treat heart burn

How to use DOMPER® TABLET

- How much to use

You should always take the lowest amount of DOMPER® TABLET that works for you and you should not take it for longer than is necessary. Although the amount of DOMPER® TABLET you should usually take is described below, your doctor may adjust your dose to your personal needs.

DOMPER® TABLET is most effective if taken 15-30 minutes before meals.

Adults and adolescents (12 years of age and over) weighing 35 kg or more:

children weighing 35kg or more:
Tablets: Take 1 tablet 3 to 4 times a day. Do not take more than 4 tablets per day

(40 mg/day).

Infants and children less than 12 years of age and weighing less than 35kg:

The effectiveness of DOMPER® TABLET has not been established in infants and children under 12 years of age with a body weight of <35 kg.

Your doctor may reduce the dose if you have kidney disorder.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Use DOMPER® TABLET for the duration that the doctor recommends.

Usually, the maximum treatment duration should not exceed one week for treatment of acute nausea and vomiting. If nausea and vomiting persists for longer than one week, you should consult your doctor. For other uses, the initial duration of treatment is up to 4 weeks.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may result in agitation, drowsiness, confusion and uncontrolled movements, such as irregular eye movements, or abnormal posture like twisted neck.

While you are using it

- Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using DOMPER® TABLET.

DOMPER[®] TABLET

Domperidone (10mg)

Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give DOMPER[®] TABLET to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Use with caution in patient with hepatic and renal impairment.

Side effects

Like all medicines, DOMPER[®] TABLET can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

DOMPER[®] TABLET is well tolerated.

The side effects which have been reported are as follow:

CNS: dry mouth, headache, insomnia, dizziness, thirst.

Gastrointestinal: transient mild abdominal cramps, diarrhoea.

Endocrinologic: Domperidone may elevate prolactin level, which may lead to mastalgia, galactorrhoea or gynecomastia.

Cardiovascular: edema, palpitations.

Paediatric population: extrapyramidal disorder.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of DOMPER[®] TABLET

- Storage

Store below 30°C, protect from light & moisture.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

- What it looks like

A white color tablet, one side with a

score “”, and the other side

impressed with a mark “”.

- Ingredients

- Active ingredient

Domperidone

- Inactive ingredient

Lactose

Povidone

Sodium Starch Glycolate

Magnesium Stearate

- MAL number

MAL19972516AZ

Manufacturer and Product

Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision:

12/09/2025

Serial Number

NPRA(R2/1)250911/01541