

# PARAS TABLET

Chlorzoxazone/ Paracetamol (250mg/300mg)

## What is in this leaflet

1. What PARAS TABLET is used for
2. How PARAS TABLET works
3. Before you use PARAS TABLET
4. How to use PARAS TABLET
5. While you are using it
6. Side effects
7. Storage and Disposal of PARAS TABLET
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What PARAS TABLET is used for

PARAS TABLET is indicated for as supplementary measure such as rest and physical therapy, for relief of pain and muscle spasm associated with acute painful musculoskeletal conditions.

## How PARAS TABLET works

Paracetamol can help in reduce pain and gives anti-inflammatory effect. Chlorzoxanone acts on specific area of the brain and involved in producing and maintaining skeletal muscle spasm.

## Before you use PARAS TABLET

### -When you must not use it

Do not take PARAS TABLET if you have known hypersensitivity to this product or have shown hypersensitivity to any component of this product.

### -Before you start to use it

Tell your doctor or pharmacist if you are allergic to any medicines or if you have any other medical conditions.

### -Taking other medicines

Tell your doctor or pharmacist if you are taking any other medications. PARAS TABLET can interact with antipsychotics drug, anti-anxiety agents or other CNS depressants. You must also inform them about all the medicines you are taking, including those without a prescription.

## How to use PARAS TABLET

### -How much to use

The usual adult dose is 1-2 tablets four times a day. It should be dispensed on physician's prescription.

### -When to use it

Consult your doctor on the period of time to take the medication.

### -How long to use it

Consult your doctor on the duration of consumption.

### -If you forget to use it

If you miss a dose of PARAS TABLET and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

### -If you use too much (overdose)

Seek for medical attention.

## While you are using PARAS TABLET

### -Things you must do

Follow your doctor's instruction when you are taking PARAS TABLET

### -Things you must not do

Do not start taking this medicine or change the dosage without first checking with your doctor. Do not give this medicine to anyone else, even if their symptoms seem similar to yours.

### -Things to be careful of

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

PARAS TABLET should be given with care to patients with liver dysfunction or a history of allergic reactions to its components. Patients should not drive or operate machine after taking this drug. It should not be given to women who might be pregnant, pregnant and lactating.

## Side effects

Gastrointestinal bleeding may occur but rare. Drowsiness, dizziness, lightheadedness can occur. Respiratory depression can occur in high doses. Rarely, it can cause rashes, edema or anaphylactic reaction. It is suspected of causing liver damage.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [portal.bpfk.gov.my](http://portal.bpfk.gov.my) (Consumers → Reporting)

## Storage and Disposal of PARAS TABLET

### -Storage


Store below 30°C, protect from light & moisture.

### -Disposal

Dispose when you stop taking the tablets as instructed by your doctor or pharmacist or the tablets have passed their expiry date. Ask them what to do with any tablets that are left over.

## Product description

### -What it looks like

A light orange round tablet with the diameter of 13.5 mm, one side impressed with a circle and a dot in the middle“”

### -Ingredients

Active ingredient-  
Chlorzoxazone  
Paracetamol  
Inactive ingredients-  
Starch  
Magnesium Stearate  
Sunset Yellow

### -MAL number

MAL19984398A

### **Manufacturer**

Y.S.P Industries (M) Sdn. Bhd.  
Lot 3,5 & 7, Jalan P/7, Section 13,  
Kawasan Perindustrian Bandar Baru Bangi,  
43000 Kajang, Selangor Darul Ehsan,  
Malaysia.

**Product Registration Holder**

Y.S.P Industries (M) Sdn. Bhd.  
Lot 3,5 & 7, Jalan P/7, Section 13,  
Kawasan Perindustrian Bandar Baru Bangi,  
43000 Kajang, Selangor Darul Ehsan,  
Malaysia.

**Date of revision :**

9 November 2015