

# TIDACT CAPSULE

Clindamycin (150 mg, 300 mg)

## What is in this leaflet

1. What TIDACT CAPSULE is used for
2. How TIDACT CAPSULE works
3. Before you use TIDACT CAPSULE
4. How to use TIDACT CAPSULE
5. While you are using it
6. Side effects
7. Storage and Disposal of TIDACT CAPSULE
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What TIDACT CAPSULE is used for

TIDACT CAPSULE is indicated for serious bacterial infections caused by Staphylococci, Streptococci (except Streptococcus faecalis), pneumococci and anaerobic bacterial infections.

## How TIDACT CAPSULE works

Clindamycin is an antibiotic which kills bacteria which are susceptible to it.

## Before you use TIDACT CAPSULE

### - When you must not use it

Do not use TIDACT CAPSULE if you are allergic towards Clindamycin or Lincomycin.

Cannot be used in newborns.

### - Before you start to use it

Please refer to doctor or pharmacist for further information.

### - Taking other medicines

Clindamycin and Erythromycin should not be administered concurrently as studies have shown antagonism in vitro.

The concurrent use of kaolin-containing anti-diarrheal preparations markedly reduces the rate of absorption of Clindamycin.

## How to use TIDACT CAPSULE

### - How much to use

Adults:

Serious infections: 150 to 300mg every 6 hours.

More severe infections: 300 to 450mg every 6 hours.

Children:

Serious infections: 8 to 16mg/kg/day divided to three or four equal doses.

More severe infections: 16 to 20mg/kg/day divided to three or four equal doses.

### - When to use it

TIDACT CAPSULE should be taken according to the time advised by your doctor or pharmacist.

### - How long to use it

Use TIDACT CAPSULE as directed by your doctor or pharmacist.

### - If you forget to use it

If you miss a dose of TIDACT CAPSULE, take it as soon as possible. If it is approaching time for the next dose, skip the missed dose and go back to your regular dosing schedule. Do not double doses.

### - If you use too much (overdose)

Seek for medical attention if you have taken too much of TIDACT CAPSULE.

## While you are using it

### - Things you must do

TIDACT CAPSULE should be used with caution in individuals with a history of gastrointestinal disease, particularly colitis.

### - Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor.

### - Things to be careful of

Safety for use in pregnancy has not been established. If pregnant or become pregnant or breast-feeding during treatment, should not take TIDACT CAPSULE unless as advised by doctor.

## Side effects

Clindamycin may cause severe and persistent diarrhea, nausea, vomiting, abdominal cramps, and abnormality of taste. Severe pseudomembranous colitis occurred in some patients and occasionally fatal. Hypersensitivity reaction including skin rashes and urticaria may occur, and transient leucopenia and eosinophilia, abnormalities of liver function tests, and jaundice have been reported. Agranulocytosis, thrombocytopenia, and erythema multiforme have been observed.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my)

[Consumers → Reporting Side Effects to Medicine (ConSERF) or Vaccines (AEFI)]

## Storage and Disposal of TIDACT CAPSULE

### - Storage

Keep in a tight container. Store below 30°C, protect from light & moisture.

### - Disposal

TIDACT CAPSULE should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required.

## Product description

### - What it looks like

TIDACT CAPSULE:

A violet colored cap and yellow colored body hard gelatine capsule with the mark "TDC", and containing white powder.

Tidact Capsule 300mg:

A size #0 light blue cap and body hard YSP gelatin capsule with the mark "TDC<sub>3</sub>" both cap and body, and containing white powder.

### - Ingredients

Active ingredient:  
Clindamycin hydrochloride

Inactive ingredients:  
Cellulose  
Magnesium Stearate  
Lactose

### - MAL number

150 mg: MAL19970979AZ

300 mg: MAL05041949AZ

## Manufacturer and Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.  
Lot 3, 5 & 7, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor Darul Ehsan, Malaysia.

## Date of revision

20/01/2021

**Serial Number**

NPRA(R2/1)210215/623