

COPASTIN FILM COATED TABLET

Cloperastine Hydrochloride 10mg

What is in this leaflet

This leaflet contains information below:

1. What COPASTIN TABLET is used for
2. How COPASTIN TABLET works
3. Before you use COPASTIN TABLET
4. How to take COPASTIN TABLET
5. While you are using COPASTIN TABLET
6. Side effects
7. Storage and disposal of COPASTIN TABLET
8. Product description
9. Manufacturer and product registration holder
10. Date of revision

What COPASTIN TABLET is used for

COPASTIN TABLET is used in the treatment of unproductive cough (dry cough).

How COPASTIN TABLET works

COPASTIN TABLET is a non-narcotic antitussive agent which suppresses unproductive cough. It works by suppressing the cough reflex by direct effect on the cough centre in the brain. It exhibits the combination of a strong central antitussive effect, papaverine-like action (for treatment of heart and brain spasm) and mild antihistaminic action.

Before you use COPASTIN TABLET

When you must not take it

You must not take COPASTIN Tablet if you are allergic to Cloperastine Hydrochloride.

Before you start to take it

Consult your doctor and pharmacist regarding the consumption of COPASTIN TABLET if you are pregnant or breastfeeding. Tell your doctor or pharmacist if you are allergic to any other medicines or you have any other health related problems.

Taking other medicines

No harmful effect reported on the concurrent use of COPASTIN TABLET with other drugs. For safety purposes, inform doctor or pharmacist about all the medicines you are taking, including those you have bought without a prescription.

How to take COPASTIN TABLET

How much to take

Adults: one to two tablets three times daily. Tablet should be taken according to the dose prescribed by your doctor or pharmacist.

When to take it

COPASTIN TABLET should be taken according to the time advised by your doctor or pharmacist.

How long to take it

COPASTIN Tablet should be taken regularly and as long as your doctor advised you to do so.

If you forgot to take it

If you miss a dose of COPASTIN TABLET and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

If you take too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause severe drowsiness, depression, confusion, tiredness and nausea.

While you are using COPASTIN TABLET

Things you must do

Follow your doctor's instruction when you are taking COPASTIN TABLET.

Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor.

Do not give COPASTIN TABLET to anyone else, even if they have the same condition as you.

Things to be careful of

COPASTIN TABLET should only be used in pregnant and nursing women when clearly needed.

Extra caution needs to be taken when used in conditions characterized by production of bronchial secretions, such as cystic fibrosis (a genetic disorder

disease that affecting the exocrine glands) as these may result in sputum retention and the development of pneumonia (inflammatory conditions of lungs).

Side effects

COPASTIN TABLET may cause anorexia, nausea, dry mouth and drowsiness. Please tell your doctor or pharmacist if you experience any other side effects.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of COPASTIN TABLET

Storage

Store below 30°C, protect from light & moisture. Keep in a tight container.

Disposal

COPASTIN TABLET should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required. Medicines should not be disposed of via wastewater or household waste. These measures will help to protect the environment.

Product description

What it looks like

COPASTIN FILM COATED TABLET is a faint green colored film coated round tablet, one side impressed with a "C".

Ingredients

Active ingredient-
Cloperastine Hydrochloride

Inactive ingredients-
Lactose
Carboxymethylcellulose Calcium
Starch
Magnesium Stearate
Neusilin
Opandry II Green
Purified water

MAL number

MAL19973698AZ

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Date of revision :

13/10/2025

Serial Number:

BPFK(R4/1)030415/00056