

TREN CAPSULE 250MG

Tranexamic Acid (250 mg)

What is in this leaflet

1. What TREN CAPSULE is used for.
2. How TREN CAPSULE works.
3. Before you use TREN CAPSULE.
4. How to use TREN CAPSULE
5. While you are using it.
6. Side effects.
7. Storage and disposal of TREN CAPSULE
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What TREN CAPSULE is used for

TREN CAPSULE is used to treat or prevent severe bleeding.

How TREN CAPSULE works

TREN CAPSULE contains an active ingredient call tranexamic acid. Tranexamic acid is an antifibrinolytic agent which regulates the balance between blood coagulation and bleeding.

Before you use TREN CAPSULE

- When you must not use it
Do not take TREN CAPSULE if you have allergy towards any components in TREN CAPSULE. Avoid taking TREN CAPSULE if you are pregnant or breastfeeding. Patients with history or risk of thrombosis (eg. stroke, heart attack, deep vein thrombosis etc.) should not take TREN CAPSULE without consultation with the doctor.
- Before you start to use it
Tell your doctor or pharmacist if you are allergic to any medicines or if you have any other health related problems (especially renal diseases and thrombotic diseases).
- Taking other medicines
Seek for your doctor or pharmacist advice if you intended to use TREN CAPSULE together with other medication, such as oral contraceptives, medication consists of Factor IX complex. Besides, you must also inform them about all the medicines you are taking, including those without a prescription.

How to use TREN CAPSULE

- How much to take
Usual dose is 4 to 6 capsules, two to three times daily.

- When to use it
TREN CAPSULE should be taken upon instruction given by the doctor or pharmacist.
 - How long to use it
TREN CAPSULE should be taken as your doctor advised you to do so.
 - If you forgot to use it
If you miss a dose of TREN CAPSULE, and you are taking it regularly, take it as soon as possible. If it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.
 - If you use too much (overdose)
Seek medical attention immediately. Symptoms of over dosage may be nausea, vomiting and orthostatic hypotension (a temporary lowering of blood pressure, usually related to suddenly standing up).
- ## While you are using TREN CAPSULE
- Things you must do
Follow your doctor's instruction when you are taking TREN CAPSULE.
 - Things you must not do
Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give this medication to anyone else, even if they have the same condition as you.
 - Things to be careful of
If you wish to take medication(s) other than TREN CAPSULE, you should consult with doctor or pharmacist. If you experience disturbances of color vision during course of treatment should promptly discontinue TREN CAPSULE and consult your doctor immediately.

Side effects

Common side effects of TREN CAPSULE are anorexia (lack or loss appetite for food), hypnosis (half-

consciousness), pruritis (itching), erythema (abnormal redness of the skin due to local congestion), skin rash, and heartburn (burning sensation in the chest).

Gastrointestinal disturbances (such as nausea, vomiting, diarrhea) may occur.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of TREN CAPSULE

- Storage
Store at temperature below 30°C. Protect from light & moisture.
- Disposal
Dispose when you stop taking the tablets as instructed by your doctor or pharmacist or the tablets have passed their expiry date. Ask them what to do with any tablets that are left over.

Product description

- What it looks like
A size # 2 orange cap, yellow body hard gelatin capsule.
- Ingredients
 - Active Ingredient-
Tranexamic acid
 - Inactive Ingredients-
Aerosil
Magnesium Stearate
Lactose
Talc
- MAL no.
MAL 19991436AZ

Manufacturer & Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia

Date of revision:

17/12/2025

Serial Number:

BPFK(R4/1)070715/00178