

UROSIN FILM COATED TABLET

Atenolol 50mg & 100mg (potency)

What is in this leaflet

This leaflet contains information below:

1. What UROSIN FILM COATED TABLET is used for
2. How UROSIN FILM COATED TABLET works
3. Before you use UROSIN FILM COATED TABLET
4. How to use UROSIN FILM COATED TABLET
5. While you are using UROSIN FILM COATED TABLET
6. Side effects
7. Storage and disposal of UROSIN FILM COATED TABLET
8. Product description
9. Manufacturer & Product Registration Holder
10. Date of revision

What UROSIN FILM COATED TABLET is used for

For the treatment of hypertension and angina pectoris.

How UROSIN FILM COATED TABLET works

UROSIN known as beta-blocker, are medication that reduces your blood pressure. UROSIN works by blocking the effects of the hormone epinephrine, also known as adrenaline. When you take UROSIN, the heart beats more slowly and with less force, thereby reducing blood pressure. UROSIN also help blood vessels open up to improve blood flow.

Before you use UROSIN FILM COATED TABLET

When you must not use it

When you are allergic to atenolol or you have asthma as well as any history of obstructive airway diseases. Please consult your doctor if you have other heart problem.

Before you start to use it

Tell your doctor if you are using any other medicines.

Taking other medicines

Salbutamol and isoprenaline can be used together with atenolol but appropriate care should be taken. Care is needed when β -blockers are combined with class I antiarrhythmic agents, (e.g. disopyramide) or calcium antagonists of the verapamil type.

How to use UROSIN FILM COATED TABLET

How much to use

For hypertension, 50-100mg per day alone or in conjunction with other antihypertensive agents. Angina pectoris: 50-100mg once a day or in divided doses for most patients. Increasing the dosage more than 100mg is unlikely to produce more beneficial effects.

When to use it

UROSIN Film Coated Tablet should be taken according to doctor's instruction.

How long to use it

UROSIN Film Coated Tablet should be taken according to doctor's advice.

If you forget to use it

If you miss a dose of UROSIN and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is nearing time for the next dose, do not double the dose to catch up, unless advised by your health care provider. Do not take 2 doses at once.

If you use too much (overdose)

Quickly seek for medical attention if you think that you or anyone else has taken too much UROSIN.

While you are using it

Things you must do

Take the tablet according to doctor's instruction.

Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction of your doctor. Do not give UROSIN to anyone else, even if they have the same condition as you.

Things to be careful of

Should be used with caution in patient with bronchospastic disease, hyperthyroidism and hypoglycemia. This product should be avoided in pregnant and nursing women.

Side effects

Bradycardia, cold extremities, postural hypotension, dizziness, vertigo, nausea, diarrhea, fatigue, skin rash, and dry eyes.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug

Reaction Monitoring by visiting the website nptra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of UROSIN FILM COATED TABLET

Storage

Keep in a tight container. Store below 30°C, protect from light & moisture.

Disposal

If your doctor or pharmacist tells you to stop taking UROSIN or the tablet has passed the expiry date, ask your doctor or pharmacist what to do with any that is left over.

Product description

What it looks like

UROSIN Film Coated Tablet 50mg:

A white round film coated tablet and one side with the mark “”

UROSIN Film Coated Tablet 100mg: A white color film coated tablet, one side

impressed with the mark “”.

Ingredients

Active ingredient:-

Atenolol

Inactive ingredient:-

Hydroxypropyl Methycellulose

Silicon Dioxide

Dimethicone

Talc

Magnesium Stearate

Cellulose

Cellulose Calcium

MAL number.

MAL 20013576A (50mg)

MAL 19962580A (100mg)

Manufacturer & Product registration Holder

Y.S.P. Industries (M) Sdn. Bhd.

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision:

06/10/2025

Serial Number

NPRA(R2/1)210408/653