

URIPAX FILM COATED TABLET

Flavoxate Hydrochloride (200mg)

What is in this leaflet

1. What URIPAX FILM COATED TABLET is used for
2. How URIPAX FILM COATED TABLET works
3. Before you use URIPAX FILM COATED TABLET
4. How to use URIPAX FILM COATED TABLET
5. While you are using URIPAX FILM COATED TABLET
6. Side effects
7. Storage and disposal of URIPAX FILM COATED TABLET
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What URIPAX FILM COATED TABLET is used for

It is used to relief bladder spasms and treat bladder symptoms such as frequent or urgent urination, increase night-time urination, painful urination and urine leakage associated with inflammation of the urinary bladder, prostate gland and urethra.

How URIPAX FILM COATED TABLET works

Flavoxate Hydrochloride is a smooth-muscle relaxant. It works by relaxing the muscles in the bladder and urinary tract.

Before you use URIPAX FILM COATED TABLET

- When you must not take it
URIPAX FILM COATED TABLET should not be used in patients who have duodenal obstruction, obstructive uropathy of the lower urinary tract and gastrointestinal hemorrhage.
- Before you start to take it
Tell your doctor or pharmacist if you are allergic to any other medicines, foods, preservatives or dyes. Tell your doctor or pharmacist if you have any other health related problems.

- Taking other medicines

Seek for your doctor or pharmacist advice if you intended to use URIPAX FILM COATED TABLET with other medicine. Some medicine and URIPAX FILM COATED TABLET may interfere with each other, especially anticholinergic or medications with anticholinergic effect and central nervous system depression-producing medications.

Besides, you must also inform them about all the medicines you are taking, including those without a prescription.

How to take URIPAX FILM COATED TABLET

Always take the tablet exactly as directed by doctor or pharmacist. Check with doctor or pharmacist if you are not sure.

- How much to take

The usual dose of URIPAX FILM COATED TABLET in adults is one tablet three times a day.

Take URIPAX FILM COATED TABLET only as directed by your doctor or pharmacist. Do not change the dose unless your doctor tells you to do so.

- When to take it

URIPAX FILM COATED TABLET should be taken according to the time advised by your doctor or pharmacist. It can be taken with or without food.

- How long to take it

Take this medicine when necessary or as long as your doctor tells you to.

- If you forgot to take it

If you miss a dose, take it as soon as you remember. However, if it is nearly time for the next dose, skip the missed dose. Do not take a

double dose to make up for a forgotten dose. Do not take two doses at once.

- If you take too much (overdose)

Contact your doctor immediately. Taking too many tablets may cause symptoms of nausea, vomiting, flushing, dilated pupils, dry mouth and tongue, hot dry skin, fever, raised heartbeat, high blood pressure, loss of full control of bodily movements, drowsiness, delirium, agitation and visual hallucinations.

While you are using URIPAX FILM COATED TABLET

- Things you must do

You must tell your doctor if you think you are or you are planning to become pregnant.

- Things you must not do

Do not start taking this medicine or change the dosage without first checking with your doctor. Do not give this medicine to anyone else, even if their symptoms or conditions seem similar to yours.

- Things to be careful of

Driving and using machines
URIPAX FILM COATED TABLET may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

URIPAX FILM COATED TABLET should be taken cautiously in patients with suspected glaucoma.

Safety use during pregnancy and lactation are unknown. Avoid use in pregnancy unless there is no safer alternative.

Exercise caution when URIPAX FILM COATED TABLET is administered during breast-feeding.

URIPAX FILM COATED TABLET

Flavoxate Hydrochloride (200mg)

Side effects

No such experience has been reported to date with URIPAX FILM COATED TABLET, but it is considered likely to have drowsiness, dizziness, headache, mental confusion (especially in the elderly), nervousness, loss of balance, fatigue, fever, blurred vision, raised heartbeat, nausea, vomiting, dry mouth, difficulty swallowing, indigestion, hypersensitivity reaction, painful urination, rashes.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website <http://npra.moh.gov.my/> (Public → Reporting Medicinal Problems / Side Effects / AEFI)

Storage and Disposal of URIPAX FILM COATED TABLET

- Storage
Store at temperature below 30°C. Protect from light and moisture. Keep out of the reach and sight of children.
- Disposal
URIPAX FILM COATED TABLET should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Return any unused or expired medicine to your doctor, pharmacist or health care professional for safe disposal

Product description

- What it looks like
A slight yellow film coated round tablet.
- Ingredients
 - Active ingredient
Flavoxate Hydrochloride
 - Inactive ingredients
Lactose
Cellulose
Magnesium Stearate
Magnesium Aluminium Silicate
Sodium Starch Glycolate

Polyethylene Glycol
Dimethicone
Titanium Dioxide

MAL no.

MAL20021159AZ

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor Darul
Ehsan, Malaysia.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi,
43000 Kajang, Selangor Darul
Ehsan,
Malaysia.

Date of revision:

19/7/2017

Serial Number:

NPRA (Rx/x)xxxxxx/xxxxx