




Item Code	: 203909 - 06 draft 19 Aug 2025
PM Code	: L.P.- IBSs 100mL
Measurement(mm) ± 2mm	: 137(L) x 50(W)
Colour of Artwork	: P. 3015 C K
Source of Artwork	: MAC 2 / CC

W
50mm
(±2mm)

YSP Ibuprofen

suspension 20 mg/mL 100 mL

Analgesic, Antipyretic

Orange Flavour

Alcohol Free

Relieves

- Fever & pain
- Muscle & joint inflammation



Scan QR code to access package insert

Each mL contains:
Ibuprofen 20 mg
Sodium Benzoate 1.2 mg (as preservative)

Indications:
Relief of fever, pain, muscle and joint inflammation.

Route of Administration:
For Oral Administration

Dosage and Administration:

Age	Dose	Frequency
10 - 12 years	15 mL	3 times daily
7 - 9 years	10 mL	3 times daily
4 - 6 years	7.5 mL	3 times daily
1 - 3 years	5 mL	3 times daily
6 - 12 months	2.5 mL	3-4 times daily
< 6 months	May be given on the advice of a doctor	

Shake well before use

Keep in a tight container.
Store at temperature below 30°C.
Protect from light and moisture.
Keep medicine out of reach of children.
Jauhkan daripada kanak-kanak.

Not recommended for Phenylketonurics.

MAL 04125482AZ

UBAT TERKAWAL
CONTROLLED MEDICINE

Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P77, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3679
Product Info: 1 800 88 3679
A Subsidiary of **Y.S.P.SAH®**



9 557892 105517

Date of opening

Batch no.

Mfg. date

Exp. date

L
137mm (±2mm)

Item Code	: 203539 - 04 draft 19 Aug 2025
PM Code	: L.P.- IBSs 4L
Measurement(mm) ± 2mm	: 237(L) x 135(W)
Colour of Artwork	: C M Y K
Source of Artwork	: MAC 2 / CC

L
237mm (±2mm)

WARNINGS

RISK OF GI ULCERATION, BLEEDING AND PERFORATION WITH NSAID

Serious GI toxicity such as bleeding, ulceration and perforation can occur at any time, with or without warning symptoms, in patients treated with NSAID therapy. Although minor upper GI problems (eg. dyspepsia) are common, usually developing early in therapy, prescribers should remain alert for ulceration and bleeding in patients treated with NSAIDs even in the absence of previous GI tract symptoms.

Studies to date have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Patients with prior history of serious adverse events and other risk factors associated with peptic ulcer disease (eg. alcoholism, smoking, corticosteroid therapy) are at increased risk. Elderly or debilitated patients seem to tolerate ulceration or bleeding less than other individuals and account for most spontaneous reports for fatal GI events.

Drug Interactions:

In therapeutic doses, no evidence of clinically significant interactions with other commonly used drugs has so far been observed. However, caution should be exercised in patients receiving oral anticoagulants, thiazide diuretics, ACE inhibitors, β blockers, methotrexate, lithium, other pain relievers containing aspirin, ibuprofen or other NSAIDs.

Side Effect (s) / Adverse Reaction(s):

Adverse reactions are usually mild and transient: Stomach discomfort, nausea, stomach ulcer, rashes or worsening of asthma, liver and kidney problems, dizziness, headache and hearing disturbance.

Symptoms And Treatment For Overdosage, And Antidote(s):

Gastric lavage may be performed and if necessary, blood electrolytes corrected.

Shelf-life:

3 years from date of manufacturer.

Product Description and Packing(s):

An orange color suspension with orange flavour.

Plastic bottle of 60ml, 100ml and 120ml.

Plastic bottle of 4L (for export only).



Scan QR code to
access package insert

KEEP IN A TIGHT CONTAINER.
STORE AT TEMPERATURE BELOW 30°C.
PROTECT FROM LIGHT AND MOISTURE.
KEEP MEDICINE OUT OF REACH OF CHILDREN.
JAUHKAN DARIPADA KANAK-KANAK.

UBAT TERKAWAL
CONTROLLED MEDICINE

BATCH NO.:

MFG DATE:

EXP DATE:

xxxxx Printing Code xxxxx

YSP Ibuprofen suspension 20mg/ml 4L

Analgesic, Antipyretic

Relieves

- Fever & pain
- Muscle & joint inflammation

Alcohol Free
Orange Flavour

MAL 04125482AZ

Each ml contains:

Ingredient(s) / Bahan Aktif:

Ibuprofen 20mg

Sodium Benzoate 1.2mg (as preservative)

Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,

43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679

A Subsidiary of **Y.S.P. SAH®**

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):

Ibuprofen is a nonsteroidal analgesic, antipyretic and anti-inflammatory agent and is effective in the relief of inflammation, pain and fever. Nonsteroidal anti-inflammatory drugs inhibit the activity of the enzyme cyclo-oxygenase, resulting in decreased formation of precursors of prostaglandins and thromboxanes from arachidonic acid.

Chemical group: A propionic acid derivative.

Half life - Elimination: 1.8 - 2 hours.

Onset of action: Pain: 0.5 hour.

Time to peak effect: Fever: 2 - 4 hours.

Duration of action -

Fever: 5mg/kg dose - 6 hours.

10mg/kg dose - 8 hours or more.

Pain: 4 - 6 hours.

Shake well before use.

Indication(s):

Relieve of fever, pain, muscle and joint inflammation.

Route of Administration: For Oral Administration.

Dosage And Administration:

Age	Dose	Frequency
10 - 12 years	15 ml	3 times daily
7 - 9 years	10 ml	3 times daily
4 - 6 years	7.5 ml	3 times daily
1 - 3 years	5 ml	3 times daily
6 - 12 months	2.5 ml	3 - 4 times daily
< 6 months	May be given on the advice of a doctor	

Contraindication(s):

Hypersensitivity to aspirin, ibuprofen or other nonsteroidal anti-inflammatory drugs. Patients with active or history of peptic ulceration. Should not be given to patients whom aspirin and other NSAIDs induce symptoms of asthma, rhinitis or urticaria.

Precaution(s) / Warning(s):

Ibuprofen suspension should be given with care to the elderly, to patients with cardiovascular disease, and in liver or renal failure.

Patients with congestive heart failure, cirrhosis, diuretic-induced volume depletion, or renal insufficiency require local synthesis of vasodilating prostaglandins to maintain renal perfusion, and therefore these patients are at greater risk of developing renal dysfunction due to NSAID-induced inhibition of renal prostaglandin synthesis.

Ibuprofen suspension should be discontinued in patients who experience blurred or diminished vision, or changes in colour vision.

Patients with collagen disease may be at increased risk of developing aseptic meningitis.

Ibuprofen suspension should be given with care to patients who have asthma or have ever suffered from asthma. If the child's symptoms (eg. earache) persist for > 3 days, consult the doctor.

Pregnancy and Lactation

Whilst no teratogenic effects have been demonstrated in animal experiments, the use of ibuprofen during pregnancy should, if possible, be avoided. In the limited studies so far available, ibuprofen appears in the breast milk in very low concentration and is unlikely to affect the breastfed infant adversely.

Not recommended for Phenylketonurics

W 135mm (±2mm)

