

Ibuprofen (20mg)

What is in this leaflet

1. What IBUPROFEN SUSPENSION is used for
2. How IBUPROFEN SUSPENSION works
3. Before you use IBUPROFEN SUSPENSION
4. How to use IBUPROFEN SUSPENSION
5. While you are using it
6. Side effects
7. Storage and Disposal of IBUPROFEN SUSPENSION
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What IBUPROFEN SUSPENSION is used for

Relieve of fever, pain, muscle and joint inflammation.

How IBUPROFEN SUSPENSION works

Ibuprofen is a nonsteroidal analgesic, antipyretic and anti-inflammatory agent. Nonsteroidal anti-inflammatory drugs inhibit the activity of the enzyme cyclooxygenase, resulting in decreased formation of precursors of prostaglandins and thromboxanes from arachidonic acid.

Before you use IBUPROFEN SUSPENSION**- When you must not use it**

Do not take this medicine if:

- you are allergic to ibuprofen or any other ingredients of this medicine (listed in Product Description)

If you are hypersensitivity to aspirin, ibuprofen or other nonsteroidal anti-inflammatory drugs. Patients with active or history of peptic ulceration. Should not be given to patients whom aspirin and other NSAIDs induce symptoms of asthma, rhinitis or urticaria.

- Before you start to use it

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breastfeeding before using any medicine.

Tell your doctor if you are allergic to any other medicines or you have any other health related problems

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Caution should be exercised in patients receiving oral anticoagulants, thiazide diuretics, ACE inhibitors, β blockers, methotrexate, lithium, and other pain relievers containing aspirin, ibuprofen or other NSAIDs.

How to use IBUPROFEN SUSPENSION**- How much to use**

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Use IBUPROFEN SUSPENSION for the duration that the doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause gastric lavage.

While you are using it**- Things you must do**

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using IBUPROFEN SUSPENSION.

Tell your doctor immediately if you become pregnant while using this medication.

Take IBUPROFEN SUSPENSION only with a doctor's prescription.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give IBUPROFEN SUSPENSION to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

IBUPROFEN SUSPENSION should be given with care to the elderly, to patients with cardiovascular disease, liver or renal and heart failure, congestive heart failure, cirrhosis, asthma, fluid retention and gastrointestinal problems.

IBUPROFEN SUSPENSION should be discontinued in patients who experience blurred or diminished vision, or changes in colour vision.

Side effects

Like all medicines, IBUPROFEN SUSPENSION can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

Adverse reactions are usually mild and transient: Stomach discomfort, nausea, stomach ulcer, rashes or worsening of asthma, liver and kidney problems, dizziness, headache and hearing disturbance.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of IBUPROFEN SUSPENSION

- Storage

Store below 30°C, protect from light & moisture.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Orange colour suspension

- Ingredients

- Active ingredient(s)
Ibuprofen

- Inactive ingredients
Glycerin
Propylene Glycol
Cellulose Sodium
Silicon Dioxide
Sunset Yellow
Tween

Aspartame
Citric Acid
Sodium Benzoate
Orange Powder

- MAL number(s):

MAL04125482AZ

Manufacturer & Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian bandar Baru Bangi, 43000 Kajang Selangor, Malaysia.

Date of revision

25/08/2025

Serial Number:

NPRA(R2/2)230505/1059