

CEPHANMYCIN CAPSULE

Cephalexin (500mg)

What is in this leaflet

1. What CEPHANMYCIN CAPSULE is used for.
2. How CEPHANMYCIN CAPSULE works.
3. Before you use CEPHANMYCIN CAPSULE.
4. How to use CEPHANMYCIN CAPSULE.
5. While you are using it
6. Side effects
7. Storage and disposal of CEPHANMYCIN CAPSULE.
8. Product Description.
9. Manufacturer and Product Registration Holder
10. Date of Revision

What CEPHANMYCIN CAPSULE is used for

Cephalexin is used to treat infections caused by bacteria, including respiratory infections, ear infections, skin infections, urinary tract infections and other infections.

How CEPHANMYCIN CAPSULE works

It works by interfering with the bacteria's cell wall formation, causing it to rupture, and killing the bacteria.

Before you use CEPHANMYCIN CAPSULE

-When you must not use it

CEPHANMYCIN CAPSULE should not be used in individuals with known allergy to cephalexin, cephalosporin group of antibiotics or any other ingredients.

-Before you start to use it

Tell your doctor if you are allergic to any other medicines such as penicillin or you have any other health related problems especially history of gastrointestinal disease.

-Taking other medicines

Seek for your doctor or pharmacist advice if you intended to use CEPHANMYCIN CAPSULE together with other medication especially metformin (antidiabetic) and probenecid (medicine for gout). Besides, you must also inform them about all the medicines you are taking, including those without a prescription.

How to use CEPHANMYCIN CAPSULE

-How much to use

Throat infection, skin infection, and uncomplicated bladder inflammation (cystitis): 2 capsules every 12 hours.

Cystitis therapy should be given only to people over 15 years of age.

For more severe infections or those caused by less susceptible organism, larger doses up to 8 capsules daily may be needed.

-When to use it

Can be taken with or without meals.

-How long to use it

Cystitis therapy should continue for 7 to 14 days. CEPHANMYCIN CAPSULE should be taken as your doctor advised you to do so.

-If you forget to use it

If you miss a dose of CEPHANMYCIN CAPSULE and you are taking it regularly, take it as soon as possible. If it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

-If you use too much (overdose)

Symptoms of oral overdose may include nausea, vomiting, stomach distress, diarrhea, and blood in the urine. Seek for medical attention.

While you are using CEPHANMYCIN CAPSULE

-Things you must do

Follow your doctor's instruction when you are taking CEPHANMYCIN CAPSULE.

-Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give CEPHANMYCIN CAPSULE to anyone else, even if they have the same condition as you.

-Things to be careful of

Safety used in pregnancy and lactation have not been established. Cephanmycin should be used during pregnancy only if clearly needed. Cephalexin is secreted in breast milk but in low concentration. Caution should be exercised when Cephanmycin is used by nursing woman.

Side effects

Some of the side effects are: Nausea and vomiting, diarrhea, indigestion, abdominal pain, hypersensitivity reaction (rash, hives, swelling and rarely, life-threatening skin

reactions), genital and anal itching, agitation, confusion, hallucination (hear, smell, taste or feel things that do not exist), joint inflammation, reversible kidney inflammation, decrease in platelet and white blood cells.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of CEPHANMYCIN CAPSULE

-Storage

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

-Disposal

CEPHANMYCIN CAPSULE should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required.

Product description

-What it looks like

A green cap and pale green body hard gelatin capsule, both cap and body  imprinted with “ 500 ” in black letters with containing white to off-white powder.

-Ingredients

-Active ingredient
Cephalexin

-In-active Ingredients

Colloidal Silicon Dioxide, Lactose Sodium Starch Glycolate, Magnesium Stearate, Talc

-MAL number.

MAL06081196AZ

Manufacturer

Yung Shin Pharmaceutical Ind. Co., Ltd.
Taichung Youth Factory III
No. 26, Kung Liu Rd., Tachia, Taichung, Taiwan, R.O.C.

CEPHANMYCIN CAPSULE

Cephalexin (500mg)

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7,
Section 13, Kawasan
Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor,
Malaysia.

Date of revision :

08/09/2025

Serial Number:

NPRA(R4/1)300919/00280