

Acknowledgement

Draft

Antibiotic 1.2g x 10Vials

CLAMENTIN INJECTION 1.2G

Each vial contains:

Amoxicillin (as Amoxicillin Sodium) 1g (potency)
Clavulanic Acid (as Potassium Clavulanate) 0.2g (potency)



Manufactured by:
YUNG SHIN PHARMACEUTICAL IND. CO., LTD.
TAICHUNG YOUTH FACTORY III
NO. 26, KUNG LIU RD., YOUTH INDUSTRIAL DISTRICT,
TACHIA, TAICHUNG, TAIWAN, R.O.C.

Imported and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD.
(199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679

Indication(s):

Clamentin Injection 1.2g is indicated for short-term treatment of bacterial infections at the following sites when Amoxicillin-resistant β -lactamase-producing strains are suspected as the cause:
Upper Respiratory Tract Infections (Including ENT) in particular sinusitis, otitis media, recurrent tonsillitis. These infections are often caused by *Streptococcus pneumoniae*, *Haemophilus influenzae**, *Moraxella catarrhalis** and *Streptococcus pyogenes*.

Lower Respiratory Tract Infections in particular acute exacerbations of chronic bronchitis (especially if considered severe), bronchopneumonia. These infections are often caused by *Streptococcus pneumoniae*, *Haemophilus influenzae** and *Moraxella catarrhalis**.

Genito-urinary Tract and Abdominal Infections in particular cystitis (especially when recurrent or complicated - excluding prostatitis), septic abortion, pelvic or puerperal sepsis and intra-abdominal sepsis. These infections are often caused by *Enterobacteriaceae** (mainly *Escherichia coli**), *Staphylococcus saprophyticus*, *Enterococcus species**.

Skin and Soft Tissue Infections in particular cellulitis, animal bites and severe dental abscess with spreading cellulitis. These infections are often caused by *Staphylococcus aureus**, *Streptococcus pyogenes* and *Bacteroides species**.

Prophylaxis of wound infection associated with surgical procedures in particular gastrointestinal, pelvic, major head and neck surgery and after limb amputation for infection.

* Some member of these species of bacteria produces β -lactamase, rendering them insensitive to Amoxicillin alone.

Dosage and Administration:

Please refer package insert for more details.

Not to be used in patients with known hypersensitivity to Penicillin.

Route of Administration:

For I.V. use only.

Store at temperature below 30°C. Protect from light and moisture.

Keep medicine out of reach of children. Jauhkan daripada kanak-kanak.

STle MYS-2160596-003

R.O.C. REG. NO.: 044070
DCA REG. NO.: MAL08042447AZ
BRU14091538P

BATCH NO.: STle

MFG. DATE:

EXP. DATE:

UBAT TERKAWAL CONTROLLED MEDICINE

不上亮油

120%

Antibiotic 1.2g x 10Vials

CLAMENTIN INJECTION 1.2G

Each vial contains:

Amoxicillin (as Amoxicillin Sodium) 1g (potency)
Clavulanic Acid (as Potassium Clavulanate) 0.2g (potency)



Manufactured by:
YUNG SHIN PHARMACEUTICAL IND. CO., LTD.
TAICHUNG YOUTH FACTORY III
NO. 26, KUNG LIU RD., YOUTH INDUSTRIAL DISTRICT,
TACHIA, TAICHUNG, TAIWAN, R.O.C.

Imported and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD.
(199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679

Indication(s):

Clamentin Injection 1.2g is indicated for short-term treatment of bacterial infections at the following sites when Amoxicillin-resistant β -lactamase-producing strains are suspected as the cause:
Upper Respiratory Tract Infections (Including ENT) in particular sinusitis, otitis media, recurrent tonsillitis. These infections are often caused by *Streptococcus pneumoniae*, *Haemophilus influenzae**, *Moraxella catarrhalis** and *Streptococcus pyogenes*.

Lower Respiratory Tract Infections in particular acute exacerbations of chronic bronchitis (especially if considered severe), bronchopneumonia. These infections are often caused by *Streptococcus pneumoniae*, *Haemophilus influenzae** and *Moraxella catarrhalis**.

Genito-urinary Tract and Abdominal Infections in particular cystitis (especially when recurrent or complicated - excluding prostatitis), septic abortion, pelvic or puerperal sepsis and intra-abdominal sepsis. These infections are often caused by *Enterobacteriaceae** (mainly *Escherichia coli**), *Staphylococcus saprophyticus*, *Enterococcus species**.

Skin and Soft Tissue Infections in particular cellulitis, animal bites and severe dental abscess with spreading cellulitis. These infections are often caused by *Staphylococcus aureus**, *Streptococcus pyogenes* and *Bacteroides species**.

Prophylaxis of wound infection associated with surgical procedures in particular gastrointestinal, pelvic, major head and neck surgery and after limb amputation for infection.

* Some member of these species of bacteria produces β -lactamase, rendering them insensitive to Amoxicillin alone.

Dosage and Administration:

Please refer package insert for more details.

Not to be used in patients with known hypersensitivity to Penicillin.

Route of Administration:

For I.V. use only.

Store at temperature below 30°C. Protect from light and moisture.

Keep medicine out of reach of children. Jauhkan daripada kanak-kanak.

STle MYS-2160596-003

R.O.C. REG. NO.: 044070
DCA REG. NO.: MAL08042447AZ
BRU14091538P

BATCH NO.: STle

MFG. DATE:

EXP. DATE:

UBAT TERKAWAL CONTROLLED MEDICINE

不上亮油