

TERASIN TABLET

Terazosin (2mg, 5mg)

What is in this leaflet

1. What TERASIN TABLET is used for
2. How TERASIN TABLET works
3. Before you use TERASIN TABLET
4. How to take TERASIN TABLET
5. While you are using TERASIN TABLET
6. Side effects
7. Storage and disposal of TERASIN TABLET
8. Product description
9. Manufacturer
10. Product Registration Holder
11. Date of Revision

What TERASIN TABLET is used for

TERASIN TABLET is used to treat symptoms of benign prostatic hyperplasia (BPH) which is benign increase in size of the prostate. It is also indicated for high blood pressure.

How TERASIN TABLET works

Terazosin is a chemical agent which improves urine flow by producing relaxation of smooth muscle in bladder neck and prostate. It lowers blood pressure by inhibiting increase of peripheral resistance in hypertensive patient. However, terazosin has no significant effect on prostate-specific antigen (PSA) level. PSA is a protein made by the cells of the prostate gland.

Before you use TERASIN TABLET

When you must not take it

Do not take TERASIN TABLET if you have allergy towards any components in TERASIN TABLET.

Before you start to take it

Tell your doctor or pharmacist if you are allergic to any medicines or if you have any other health related problems. Patient should be examined to rule out presence of carcinoma of the prostate before TERASIN treatment.

Taking other medicines

Seek for your doctor or pharmacist advice if you intended to use TERASIN TABLET together with other medication. Besides, you must also inform them about all the medicines you are taking, including those without a prescription.

How to take TERASIN TABLET

How much to take

Recommended starting dose: 1mg at bedtime.

Subsequent dose:

-Hypertension

The usual recommended dose is 1 to 5mg, once a day.

-Benign Prostatic Hyperplasia (BPH)

The usual recommended dose is 5 to 10mg, once a day.

When to use it

TERASIN TABLET should be taken at bedtime.

How long to use it

TERASIN TABLET should be taken as your doctor advised you to do so.

If you forgot to use it

If you miss a dose of TERASIN TABLET, take the missed dose as soon as you remember. However, if it's almost time for the next dose, skip the missed dose and continue the regular dosing schedule. Do not take a double dose to make up for a missed one.

If you take too much (overdose)

Seek medical attention immediately.

While you are using TERASIN TABLET

Things you must do

Follow your doctor's instruction when you are taking TERASIN TABLET. Advisable to measure blood pressure 2-3 hours after dosing to evaluate the symptoms such as dizziness or palpitations. If TERASIN TABLET was discontinued for several days or longer, therapy should be restart with initial dosing regimen.

Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give this medication to anyone else, even if they have the same condition as you.

Things to be careful of

If you are taking medications other than TERASIN TABLET, please consult with your doctor or pharmacist. If you are required to drive or operate heavy machine, please be careful of syncope (fainting or passing out) associate with first few dose of TERASIN TABLET.

Side effects

Possible side effects of TERASIN TABLET are low blood pressure (hypotension), weakness (asthenia), blurred vision, dizziness, nasal congestion, nausea, peripheral edema(swelling of the feet, ankles and legs), palpitation (abnormality of heartbeat), drowsiness(somnolence) and inability to develop or maintain an erection of the penis during sexual activity (impotence). The adverse reactions are usually mild or moderate in intensity but sometimes were serious enough to interrupt treatment. If you encounter any other side effect that is not listed here, please refer to your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of TERASIN TABLET

Storage

Store below 30°C, protect from light & moisture.

Disposal

TERASIN TABLET should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required. Medicines should not be disposed of via wastewater or household waste. These measures will help to protect the environment.

Product description

What it looks like

Terasin Tablet 2mg:

A round yellow tablet and both sides impressed with circle in the middle with cross by the side



Terasin Tablet 5mg:

A round brownish red tablet and both sides impressed with circle in the middle with cross by the side



Ingredients

Active Ingredient-
Terazosin

Inactive Ingredients-

Lactose

Sodium Starch Glycolate

Povidone

Magnesium Stearate

Talc

(For Terasin 5mg only):

Iron Oxide Red

Sunset Yellow Lake

(For Terasin 2mg only):

Quinoline Yellow

Sunset Yellow

MAL numbers

Terasin 2mg - MAL07050027AZ

Terasin 5mg - MAL07050026AZ

**Manufacturer and Product Registration
Holder**

Y.S.P Industries (M) Sdn. Bhd.

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor
Malaysia..

Date of revision :

05/08/2025

Serial Number

BPFK(R4/1)080115/00007