

CLAMENTIN FILM COATED TABLET 625MG

Amoxycillin /Clavulanic Acid (625mg)

What is in this leaflet

1. What CLAMENTIN FILM COATED TABLET 625MG is used for
2. How CLAMENTIN FILM COATED TABLET 625MG works
3. Before you use CLAMENTIN FILM COATED TABLET 625MG
4. How to use CLAMENTIN FILM COATED TABLET 625MG
5. While you are using it
6. Side effects
7. Storage and Disposal of CLAMENTIN FILM COATED TABLET 625MG
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What CLAMENTIN FILM COATED TABLET 625MG is used for

It is used for short-term treatment of bacterial infection at respiratory tract, urinary tract, skin and soft tissue, bone and joint and dental.

Your doctor may have prescribed CLAMENTIN FILM COATED TABLET 625MG for another reason. Ask your doctor if you have any questions about why CLAMENTIN FILM COATED TABLET 625MG has been prescribed for you.

How CLAMENTIN FILM COATED TABLET 625MG works
CLAMENTIN FILM COATED TABLET 625MG consists of active substance Amoxycillin/ Clavulanic Acid. These medicine work by killing the bacteria, which cause infections.

Before you use CLAMENTIN FILM COATED TABLET 625MG

- When you must not use it

Do not use CLAMENTIN FILM COATED TABLET 625MG if:

- you have had an allergic reaction to penicillin or similar types of antibiotics (such as cephalosporins) or any of the ingredients contained in CLAMENTIN FILM COATED TABLET 625MG. The ingredients are listed at the end of this leaflet. Some of the symptoms of an allergic reactions may include skin rash, itching, difficulty in breathing and swelling of the face or tongue.

- you have previously experienced liver problems after using CLAMENTIN FILM COATED TABLET 625MG or any other penicillin antibiotic.

- Before you start to use it

Tell your doctor if:

- you have ever had an allergic reaction (such as a rash) to antibiotics such as penicillin or other substances in the past.
- you are allergic to foods, dyes, preservatives or any other medicines.
- you have experienced liver problems after using CLAMENTIN FILM COATED TABLET 625MG or any other medicines.
- you have glandular fever (mononucleosis).
- you have kidney or liver problem.

Pregnancy and lactation

Please consult your doctor before start to use CLAMENTIN FILM COATED TABLET 625MG if you are pregnant, plan to become pregnant, breast-feeding or intend to breast-feed. Your doctor will have weighed the risks of you taking CLAMENTIN FILM COATED TABLET 625MG against the benefits they expect it will have for you.

CLAMENTIN FILM COATED TABLET 625MG is not recommended for children of 12 years and under.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines when taken with CLAMENTIN FILM COATED TABLET 625MG may interfere with each other. These include:

- anticoagulant therapy, medicines for blood thinning effect.
- oral contraceptives.
- probenecid, medicine for gout.
- allopurinol, medicine for gout.

Your doctor or pharmacist may have more information on medicines to be careful with or to avoid while taking medication.

How to use CLAMENTIN FILM COATED TABLET 625MG

- How much to use

Adults and children over 12 years

Mild – Moderate infections:

One tablet CLAMENTIN FILM COATED TABLET 625MG two times a day.

Your doctor might prescribe a lower dosage for you, if you have kidney or liver disease.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue using CLAMENTIN FILM COATED TABLET 625MG for as long as your doctor recommends.

Do not stop taking medication, even if you feel better after a few days, unless advised by your doctor. Your infection may not clear completely if you stop taking your medication too soon.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

If you miss a dose of CLAMENTIN FILM COATED TABLET 625MG and you are taking it regularly, take it as soon as possible. If it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause gastrointestinal symptoms and disturbance of the fluid and electrolytes balance.

While you are using it**- Things you must do**

Follow your doctor's instruction when you are taking CLAMENTIN FILM COATED TABLET 625MG.

Tell all the doctors, dentists and pharmacists treating you that you are taking CLAMENTIN FILM COATED TABLET 625MG.

Tell your doctor immediately if you become pregnant and intended to breast-feed while taking this medicine.

Ask your doctor or pharmacist for advice before taking any medicine.

If you get severe diarrhoea, tell your doctor or pharmacist immediately.

- Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give CLAMENTIN FILM COATED TABLET 625MG to anyone else, even if they have the same condition as you.

- Things to be careful of**Driving and using machine**

Be careful driving or operating machinery until you know how CLAMENTIN FILM COATED TABLET 625MG affects you.

Side effects

CLAMENTIN FILM COATED TABLET 625MG may cause gastrointestinal side effect such as diarrhea, indigestion, nausea and vomiting. It may give hematological (blood disorder) effects such as thrombocytopenia, haemolytic anemia and cause hypersensitivity reactions

such as urticarial (itching) and erythematous rashes, severe allergic reaction such as Stevens-Johnson syndrome, toxic epidermal necrolysis and exfoliative dermatitis. It also could cause hyperactivity, dizziness, headache and phlebitis (inflammation of vein).

Like all medicines, CLAMENTIN FILM COATED TABLET 625MG can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any other side effects after taking this medicine.

Stop taking CLAMENTIN FILM COATED TABLET 625MG and contact your doctor immediately if you experience any of the following:

Serious allergic reactions such as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). DRESS appears initially as flu-like symptoms with a rash on the face and then with an extended rash, high temperature and enlarged lymph nodes.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of CLAMENTIN FILM COATED TABLET**- Storage**

Store at temperature below 30°C. Protect from light and moisture.

- Disposal

CLAMENTIN FILM COATED TABLET 625MG should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your doctor and pharmacist how to dispose medicines that are no longer required.

Product description**- What it looks like**

An elliptical white film coated tablet, one side impressed with "YSP 3"

**- Ingredients**

- Active ingredient(s)
Amoxicillin Trihydrate
Clavulanate Potassium

- Inactive ingredients:
Cellulose
Magnesium Stearate
Silicon Dioxide
Sodium Starch Glycolate
Titanium Dioxide
Triethyl Citrate
Methylene Chloride

- MAL number

MAL07124630AZ

Manufacturer

Yung Shin Pharmaceutical Ind. Co., Ltd.
(Taichung Youth Factory III)
No. 26 Kung Liu Rd. Youth Industrial District, Taichia, Taichung, Taiwan, R.O.C

Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision

10/09/2020

Serial Number

NPRA(R2/1)200909/515