

LIPROCT[®] SUPPOSITORY

Lidocaine, Hydrocortisone acetate, Zinc oxide (60mg, 5mg, 400mg)

What is in this leaflet

1. What LIPROCT SUPPOSITORY is used for
2. How LIPROCT SUPPOSITORY works
3. Before you use LIPROCT SUPPOSITORY
4. How to use LIPROCT SUPPOSITORY
5. While you are using it
6. Side effects
7. Storage and disposal of LIPROCT SUPPOSITORY
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What LIPROCT SUPPOSITORY is used for

LIPROCT SUPPOSITORY is used to relieve pain, itching and discomfort around the anus (back passage) caused by certain conditions.

How LIPROCT SUPPOSITORY works

LIPROCT SUPPOSITORY contains three active ingredients which are lidocaine (local anaesthetic) that provides fast relief by numbing the area, hydrocortisone acetate (corticosteroid) that reduces swelling, redness, and itching and zinc oxide (skin protectant) which protects and soothes the irritated skin while helping it heal.

Before you use LIPROCT SUPPOSITORY

- When you must not use it

Do not use this medicine if you:

- are allergic to lidocaine, hydrocortisone acetate, zinc oxide, other amide-type local anaesthetics or any other ingredients of this medicine (listed in Product Description)
- have an untreated infection around the anal area caused by bacteria, viruses, fungi, or parasites

- Before you start to use it

Tell your doctor or pharmacist if you are allergic to any medicines or you have any other health related problems.

Check with your doctor before using this medicine if you have not been diagnosed with the cause of your rectal symptoms. Some symptoms may be caused by a more serious condition, for example malignancy.

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

Small amounts of the medicine can pass into breast milk, but it is unlikely to harm your baby when used at the recommended dose. Use only as advised by your doctor.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Tell your doctor or pharmacist if you are taking:

- medicines for an irregular heartbeat
- other medicines that contain local anaesthetics (numbing medicines)

How to use LIPROCT SUPPOSITORY

- How much to use

- Insert 1 suppository into the anus in the morning, at night, and after each bowel movement (passing stool).
- Insert the suppository just inside the anus.
- Do not use more than 5 suppositories in one day.

- When to use it

Insert one suppository morning and night and after defecation. If passing stool is painful, insert the suppository a few minutes before your bowel movement.

- How long to use it

- Treatment usually lasts 10 days to 3 weeks, or as advised by your doctor.
- If you need to use this medicine for a longer period, your doctor may advise you to stop for a short time before restarting.
- If irritation develops or does not go away, stop using the medicine and consult your doctor.

- If you forgot to use it

Use the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to use the medicine and skip the missed dose. Do not use a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Using too much of this medicine may cause dizziness, drowsiness, confusion, shaking, seizures (fits), difficulty breathing, or an irregular heartbeat.

While you are using LIPROCT SUPPOSITORY

- Things you must do

Use LIPROCT SUPPOSITORY exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using LIPROCT SUPPOSITORY.

Tell your doctor immediately if you become pregnant while using LIPROCT SUPPOSITORY.

- Things you must not do

Do not stop using LIPROCT SUPPOSITORY unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

LIPROCT® SUPPOSITORY

Lidocaine, Hydrocortisone acetate, Zinc oxide (60mg, 5mg, 400mg)

Do not give LIPROCT SUPPOSITORY to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Stop using the medicine and see your doctor if you develop irritation, worsening pain, or bleeding from the rectum.

In rare cases, the medicine may be absorbed into the body and affect the brain or nervous system, especially if large amounts are used. Seek medical help immediately if seizures (fits) occur.

Do not use this medicine for longer than recommended. Long-term or excessive use may cause thinning of the skin or other steroid-related side effects, although these are unlikely when used as directed.

Side effects

Like all medicines, LIPROCT SUPPOSITORY can cause side effects, although not everybody gets them.

Possible side effects are:

- Some people may develop an allergic skin reaction, such as redness, itching, rash, or irritation, where the medicine is used.
- In very rare cases, a severe allergic reaction may occur. Get medical help immediately if you develop difficulty breathing, swelling of the face, lips, tongue, or throat, or severe rash.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npdra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of LIPROCT SUPPOSITORY

- Storage

Store below 25°C, protect from light and moisture.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

- What it looks like

White to off-white spindle suppositories.

- Ingredients

- Active ingredient:
Lidocaine
Hydrocortisone acetate
Zinc oxide

- Inactive ingredients:
Witepsol
Aluminium Subacetate

- MAL number

MAL09011474AZ

Manufacturer and Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor, Malaysia.

Date of Revision

24/08/2026

Serial Number

NPRA(R4/1)080726/00034