

CONPANZOLE® ENTERIC COATED TABLET

Pantoprazole (20mg, 40mg)

What is in this leaflet

1. What CONPANZOLE® is used for
2. How CONPANZOLE® works
3. Before you use CONPANZOLE®
4. How to use CONPANZOLE®
5. While you are using it
6. Side Effects
7. Storage and Disposal of CONPANZOLE®
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What CONPANZOLE® is used for

CONPANZOLE® is used in combination with antibiotics for the eradication of *Helicobacter pylori* in patients with peptic ulcers. It is used also for duodenal ulcer, gastric ulcer moderate and severe cases of inflammation of the esophagus, Zollinger-Ellison-Syndrome and other hypersecretory conditions (a condition where your body secretes too many gastric acid).

How CONPANZOLE® works

CONPANZOLE® contains the active substance Pantoprazole. Pantoprazole is a selective “proton pump inhibitor”, a medicine which reduces the amount of acid produced in your stomach. It is used for treating acid-related disease of the stomach.

Before you use CONPANZOLE®

- When you must not use it

You must not take CONPANZOLE® if you are allergic to the ingredient or any other component of this medicine. You must not use this medicine if you are allergic to other proton pump inhibitors.

- Before you start to use it

Tell your doctor or pharmacist if you are allergic to Pantoprazole or any components in this product.

Tell your doctor before taking this medicine, if you are due to have a specific blood test (Chromogranin A).

- Taking other medicines

Seek advice from your doctor or pharmacist, if you intend to use CONPANZOLE® together with other medication.

Besides that, you must also inform your doctor or pharmacist about all the medicines you are taking, including those without a prescription.

How to use CONPANZOLE®

- How much to use

The daily consumption dose ranges from 20-40mg depending on different conditions. Please take according to your doctor's prescription.

- When to use it

Take the tablets 1 hour before a meal, without chewing or crushing them, and swallow them whole with some water.

- How long to use it

The duration to take CONPANZOLE® depends on your symptoms and condition. Please follow your doctor's prescription.

- If you forget to use it

Take the missed dose as soon as you remember it. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. Do not take a double dose to make up for a missed one.

- If you use too much (overdose)

Seek medical attention if you think you or anyone else may have taken too much of this supplement CONPANZOLE®.

While you are using it

- Things you must do

Follow your doctor's instruction when you are taking CONPANZOLE®.

- Things you must not do

Do not stop use this medicine or change the dosing regimen without instruction from your doctor. Do not give this medication to anyone else, even if they have the same condition as you.

- Things to be careful of

You should not take CONPANZOLE® for mild gastrointestinal problems such as indigestion. It should not be given to children less than 12 years of age unless advised by doctor.

Side effects

When you are taking this medicine, your doctor will want to monitor you (especially if you are taking it for long term). Hence, you should report any new and exceptional symptoms and circumstances whenever you see your doctor. Please tell your doctor promptly if you get any of the symptoms below:

- Rash (especially in areas exposed to the sun), possibly with pain in the joints. (Subacute Cutaneous Lupus Erythematosus, SCLE) (frequency unknown).
- Fever, extreme tiredness, pus/ blood in urine
- Involuntary muscle contractions, disorientation, convulsions, dizziness, increased heart rate
- Fracture in the hip, wrist or spine (frequency uncommon). Tell your doctor if you have osteoporosis if you have taken corticosteroids (which can increase the risk of osteoporosis).

CONPANZOLE® ENTERIC COATED TABLET

Pantoprazole (20mg, 40mg)

- Inflammation in the large bowel, that causes persistent watery diarrhea (frequency not known)
- Watery stool, stomach pain and fever that do not go away
- Anemic (pale skin, weakness, tiredness or lightheadedness), shortness of breath, a smooth tongue, nerve problems (numbness or tingling, muscle weakness and problems walking), vision loss and mental problems (depression, memory loss or behavioral changes).
- Headache, dizziness, blurred vision, diarrhea, nausea, abdominal pain and discomfort.
- Kidney problems (interstitial nephritis).
- Low levels of magnesium (hypomagnesaemia) can also lead to a reduction in potassium or calcium levels in the blood (frequency unknown).
- Severe diarrhea which may be caused by an infection (*Clostridium difficile*) in your intestines.
- Benign polyps in the stomach (frequency common).
- Proton pump inhibitors may cause vitamin B12 deficiency.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of CONPANZOLE®

- Storage

Store at temperature below 30°C. Protect from light and moisture.


- Disposal


CONPANZOLE® should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date.

Ask your pharmacist how to dispose medicines that are no longer required.

Product Description

- What it looks like

20mg: A yellow color oval shape enteric coated tablet” ”

40mg: A yellow color oval shape enteric coated tablet.” ”

- Ingredients

- Active ingredient(s)

Pantoprazole

- Inactive ingredients

Sodium Carbonate, Cellulose, Lactose, Talc, Sodium Stearyl Fumarate, Crospovidone, Polyethylene Glycol, Titanium Dioxide, Sodium Bicarbonate, Sodium Lauryl Sulfate, Opadry Yellow, Methacrylic Acid Copolymer.

- MAL number:

20mg: MAL11095002AZ

40mg: MAL11095003AZ

Manufacturer and Product

Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision

26/03/2026

Serial Number

NPRA (R2/1)210921/754