

# KASTIPRON® TABLET

Kastipron Tablet (S-Carboxymethylcysteine) is a mucolytic expectorant.

## Ingredient(s):

Each tablet contains:

S-Carboxymethylcysteine ..... 375mg

## Pharmacodynamics:

1. S-Carboxymethylcysteine possesses mucolytic and mucoregulatory properties. It has a direct effect on the mucopolysaccharides of the mucus to reduce the viscosity of the secretions, to facilitate expectoration and to clear the respiratory airways.
2. S-Carboxymethylcysteine is used in the treatment of disorders of the respiratory tract associated with excessive mucus.

## Pharmacokinetics:

1. It is very quickly absorbed in the digestive tract. Blood concentration reach a maximum after one hour and became nil 12 hours following ingestion.
2. S-Carboxymethylcysteine is mainly excreted in the urine as unchanged drug and metabolites.

## Indication(s):

Aids expectoration by reducing sputum viscosity.

## Dosage and Administration:

The usual dose for adult is 2 tablets 3 times daily after meals. A suggested dose for those 6 – 12 years is 1 tablet twice daily.

## Route of Administration:

Oral

## Contraindication(s):

Contraindicated in patients who have hypersensitivity to this product.

Contraindicated in children below two (2) years of age.

## Precaution(s) / Warning(s):

1. To be used with caution in patients with a history of peptic ulcer and to be avoided in patients with active ulceration.
2. Safety for use in pregnancy has not been established. Although animal reproductive studies have not resulted in any teratogenic effects, it should not be used in pregnant women unless benefit outweighs the risk.

## Side Effect(s) / Adverse Reaction(s):

Nausea, headache, gastric discomfort, diarrhea, gastrointestinal bleeding and skin rash have occasionally occurred with S-Carboxymethylcysteine.

## Immune System Disorders:

Anaphylactic / anaphylactoid reaction.

## Skin and Subcutaneous Tissue Disorders:

Severe cutaneous adverse reactions (SCAR) e.g. erythema multiforme, Steven-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN). In most of these cases reported at least one other drug was administered at the same time, which may have possibly enhanced the described mucocutaneous effects.

**Symptoms and Treatment for Overdosage, and Antidote(s):**

There is no specific antidote for its overdose, therefore management of the patient consists of symptomatic and supportive therapy.

**Storage Condition(s):**

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

**Shelf-Life:**

3 years from the date of manufacture.

**Product Description:**

A white to off-white color round tablet, one side impressed with a score and the other side with a mark “A”. “



**Packing(s):**

Plastic container of 200’s, 1000’s, 1200’s and 1500’s (for export only)

Blister packing of 10’s x 10 and 10’s x 100.

---

# KASTIPRON® TABLET

**Bahan Aktif:**

Setiap tablet mengandungi:

S-Carboxymethylcysteine ..... 375mg

**Indikasi:**

Mengurangkan kelikatan kahak dalam bantuan ekspektorasi.

**Dos:**

Dos biasa untuk orang dewasa adalah 2 tablet 3 kali sehari selepas makan. Untuk kanak-kanak berumur 6 – 12 tahun: 1 tablet 2 kali sehari.

---

# 痰可出錠

**成份:**

每錠中含:

S-Carboxymethylcysteine ..... 375mg

**適應症:**

減少呼吸道粘膜分泌的粘稠性。祛痰劑。

**用法、用量:**

成人每次2 錠, 每日3次, 6 - 12歲, 每次 1 錠,每日2次, 飯後服用。



Manufacturer and Product Registration Holder:  
**Y.S.P. INDUSTRIES (M) SDN. BHD.** (199001001034)  
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,  
Kawasan Perindustrian Bandar Baru Bangi,  
43000 Kajang, Selangor, Malaysia.  
Ordering Line: 1 800 88 3027  
Product Info: 1 800 88 3679