

SWECON[®] TABLET

Magnesium Aluminium Hydroxide Carbonate (Hydrotalcite) (500mg)

What is in this leaflet

1. What Swecon[®] Tablet is used for
2. How Swecon[®] Tablet works
3. Before you use Swecon[®] Tablet
4. How to use Swecon[®] Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Swecon[®] Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Swecon[®] Tablet is used for

Swecon[®] Tablet is used for gastric ulcer, inflammation of stomach lining and gastric hyperacidity.

How Swecon[®] Tablet works

Swecon[®] Tablet works by lowering the amount of acid in your stomach.

Before you use Swecon[®] Tablet

- When you must not use it
Do not take Swecon[®] Tablet if you have kidney disorder, heart disorder, diarrhea and high magnesium level.
- Before you start use it
Tell your doctor or pharmacist if you are allergic to any other medicines or you have any other health related problems.
- Taking other medicines
Do not take Swecon[®] Tablet within 1 to 2 hours after administering tetracycline. Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Swecon[®] Tablet

- How much to use
To be chewed before swallowing.
Adult: 2 tablets 3 – 4 times a day.
Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Take on an empty stomach between meals or before sleep, depending on the age and severity of the case. Use as directed by your doctor or pharmacist.

- How long to use it

Take Swecon[®] Tablet for the duration that the doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using Swecon[®] Tablet

- Things you must do
Tell all the doctors, dentists and pharmacists treating you that you are using Swecon[®] Tablet.

Tell your doctor immediately if you become pregnant while using this medication.
- Things you must not do
Do not stop taking this medicine or change the dosing regimen without instruction from your doctor.

Do not give Swecon[®] Tablet to anyone else, even if they have the same condition as you.

- Things to be careful of

This product should be used with caution in individuals with kidney impairment. Caution should be exercised when used by pregnant women.

Side effects

Occasionally: Diarrhea, soft stool, dry mouth and eating disorders.


You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Swecon[®] Tablet

- Storage
Keep out of the reach and sight of children.
Store below 30°C, protect from light & moisture.
- Disposal
Swecon[®] Tablet should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required.

Product Description

- What it looks like
A round concurve white tablet, one side is impressed with 'SWECON' and the other side is impressed a circle with a dot at the center.



- Ingredients

Active ingredients:

Magnesium Aluminium Hydroxide
Carbonate (Hydrotalcite)

Inactive ingredients:

Sodium Saccharin
Sodium Starch Glycolate
Starch
Cellulose
Magnesium Stearate
Polyethylene Glycol
Menthol
Pineapple Flavour
Milk Flavour
Povidone
Magnesium Aluminium Silicate

- MAL number(s):

MAL19972152X

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor D.E.,
Malaysia.

Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor D.E.,
Malaysia.

Date of revision

01/02/2021

Serial Number

NPRA(R4/1)010221/00005