

PORO[®] SUSPENSION 250MG/5ML (STRAWBERRY FLAVOUR)

Paracetamol (250mg/5mL)

What is in this leaflet

1. What Poro[®] Suspension is used for
2. How Poro[®] Suspension works
3. Before you use Poro[®] Suspension
4. How to use Poro[®] Suspension
5. While you are using Poro[®] Suspension
6. Side effects
7. Storage and Disposal of Poro[®] Suspension
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Poro[®] Suspension is used for

Poro[®] Suspension is used for the relief of fever and mild to moderate pain.

How Poro[®] Suspension works

Poro[®] Suspension consists of paracetamol, which is an analgesic that relieves pain and an antipyretic that reduces fever.

Before you use Poro[®] Suspension

- When you must not use it

Do not use Poro[®] Suspension if you have allergy towards paracetamol or any components in this preparation.

- Before you start to use it

Tell your doctor or pharmacist you are allergic to paracetamol or you have any other health related problem. Inform your healthcare provider if you have suffered from anemia (lack of healthy red blood cells), heart diseases, lung-related diseases, kidney or liver problems.

- Taking other medicines

Do not take other paracetamol containing medicines at the same time. Inform your doctor or pharmacist if you are taking aspirin or anticoagulants (medicines that can retard or inhibit blood clot).

Besides, you must also inform your healthcare providers about all the medicines you are taking, including those without a prescription.

How to use Poro[®] Suspension

- How much to use

Children 1 – 5 years:

2.5mL – 5mL (1/2– 1 teaspoon)

Children 6 – 12 years:

5mL – 10mL (1 – 2 teaspoon)

Doses may be repeated every 4 to 6 hours when necessary (Maximum of 4 doses in 24 hours). Do not exceed the prescribed dose.

- When to use it

Poro[®] Suspension should be taken according to the time advised by your doctor or pharmacist. It can be taken with or without food.

- How long to use it

Use Poro[®] Suspension as directed by your doctor or pharmacist.

If fever persists for more than 3 days or recurs, or if pain continues for more than 5 days, consult your doctor immediately.

- If you forget to use it

If you have missed a dose of Poro[®] Suspension, take it as soon as possible. If it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention. Taking too much may cause: nausea, vomiting, low blood pressure and sweating.

While you are using Poro[®]

Suspension

- Things you must do

Follow your doctor's instruction when you are taking Poro[®] Suspension.

- Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give Poro[®] Suspension to anyone else, even if they have the same condition as you.

WARNING

This preparation contains
PARACETAMOL.

Do not take any other paracetamol containing medicines at the same time.

- Things to be careful of

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

If fever persists for more than 3 days or recurs, or if pain continues for more than 5 days, consult a doctor immediately.

Side effects

Side effects of paracetamol are usually mild, though events of abnormal blood counts have been reported. Besides, skin rashes and other allergic reactions occur occasionally.

Skin allergic reactions include skin rashes, angioedema (swelling of deeper layer of skin caused by build-up of fluid), Steven-Johnson Syndrome (skin disorder manifested by painful rash that spreads and blisters) or Toxic Epidermal Necrolysis (skin disorder

PORO[®] SUSPENSION 250MG/5ML (STRAWBERRY FLAVOUR)

Paracetamol (250mg/5mL)

characterized by widespread reddening and blistering of the skin which might spread to eye, mouth and genital areas).

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of PORO[®] SUSPENSION

- Storage

Keep in an airtight container. Store at temperature below 30°C. Protect from light and moisture. Keep medicine out reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Pink color, strawberry-flavoured suspension.

- Ingredients

Active ingredient:
Paracetamol

Inactive ingredient(s):

Ponceau 4R
Propyl Paraben
Tween 80
Strawberry Oil
Methyl Paraben
Aerosil
Glycerin
Citric Acid
Purified Water
Sodium Saccharine
Carboxymethylcellulose Sodium
Sugar

Sodium Citrate

- MAL number:

MAL19993051XZ

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision

02/05/2025

Serial Number

NPRA(R4/1)120820/00118