

Homecare® DAVMORID® FILM COATED TABLET

Ingredient(s):

Each tablet contains:

Diosmin	450mg
Hesperidin	50mg

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetic):

The efficacy of Davmorid is accounted for by its specific action on the principle elements of venous disease. It reinforces venous tone by prolonging the activity of parietal noradrenaline. Thus, Davmorid decreases venous capacitance, venous distensibility and venous emptying time. It protects the microcirculation by fighting the microcirculation-damaging process; it combats venous inflammation by decreasing leukocyte activation and as a consequence, by inhibiting the release of inflammatory mediators, principally free radicals and prostaglandins. Thus, Davmorid normalizes capillary permeability and strengthens capillary resistance. It acts on the lymphatic system: it improves lymphatic drainage by increasing lymph flow and lymph oncotic pressure. This action on the lymphatic system associated with a phlebotonic and vasculoprotective effect, explains the activity of Davmorid on CVI-associated edema. It also significantly reduces the frequency, severity and duration of acute hemorrhoidal attacks.

Oral administration of substance containing 14C-Diosmine to man showed the following:

- 1) Excretion is mainly in faecal, urine excretion being 14% of the administered quantity.
- 2) The substance is extensively metabolised, and metabolisation is detected by the presence of different phenol acids in the urine.

Indication(s):

Treatment of organic and functional chronic venous insufficiency of the lower limbs with the following symptoms: heavy legs, pain and nocturnal cramps. Treatment of chronic hemorrhoids and acute hemorrhoidal attacks.

Dosage and Administration:

Usual dose in chronic venous insufficiency: 2 tablets daily in two divided doses morning and evening at mealtimes.

In acute hemorrhoidal attacks: 6 tablets daily in 2 divided doses, for 4 days and then 4 tablets daily in 2 divided doses for 3 days then 2 tablets daily.

In chronic hemorrhoidal disease: 2 tablets daily.

Tablets are to be taken after meals.

Side Effect(s) / Adverse Reaction(s):

Some cases of minor gastrointestinal and neurovegetative disorders have been reported, which never required cessation of treatment.

Contraindications:

Hypersensitivity to Diosmin or Hesperidin, or to any of the ingredients.

Precaution(s) / Warning(s):

Use in pregnancy:

Experimental studies have not shown any teratogenic effect in animals. In human beings, no harmful effect has so far been reported. Use in lactation: In the absence of data concerning the diffusion into breast milk, breastfeeding is not recommended during treatment.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description and Packing(s):

A light brown color film coated elliptical tablet. "



Plastic bottle of 100's. (for export only)

Blister packing of 10's X 3 and 10's X 50.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679