HOMECARE BAROSCON DOUBLE ACTION SUSPENSION

Sodium Alginate 500mg, Calcium Carbonate 325mg, Sodium Bicarbonate 213mg

What is in this leaflet

1. What HOMECARE BAROSCON DOUBLE ACTION SUSPENSION is used for
2. How HOMECARE BAROSCON DOUBLE ACTION SUSPENSION works
3. Before you use HOMECARE BAROSCON DOUBLE ACTION SUSPENSION
4. How to use HOMECARE BAROSCON DOUBLE ACTION SUSPENSION
5. While you are using it
6. Side effects
7. Storage and Disposal of HOMECARE BAROSCON DOUBLE ACTION SUSPENSION
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What HOMECARE BAROSCON DOUBLE ACTION SUSPENSION is used for

Treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, and for symptoms of excess stomach acid (hyperacidity).

How HOMECARE BAROSCON DOUBLE ACTION SUSPENSION works

HOMECARE BAROSCON DOUBLE ACTION SUSPENSION is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate. It works in two ways:

1) Neutralising excess stomach acid to relieve the pain and discomfort

2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest

Before you use HOMECARE BAROSCON DOUBLE ACTION SUSPENSION

- *When you must not use it*

Do not take this medicine if you have an allergy to any of the ingredients including the esters of hydroxybenzoates.

- *Pregnancy and lactation*

There is no significant adverse effects of the product known on the course of pregnancy or on the health of the foetus/ new-born child.

- *Before you start to use it*

Tell your doctor if you are allergic to any medicines or if you have any other medical condition.

- *Taking other medicines*

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use HOMECARE BAROSCON DOUBLE ACTION SUSPENSION

- *How much to use*

Adults and children aged 12 and above: 1 to 2 sachets after meals and before bedtime.

If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- *When to use it*

Use as directed by your doctor or pharmacist.

- *How long to use it*

HOMECARE BAROSCON DOUBLE ACTION may be used after meals and before bedtime. You may stop taking this product when you have no more symptoms of gastro-oesophageal reflux.

- *If you forget to use it*

If you miss dose of HOMECARE BAROSCON DOUBLE ACTION and you are taking it regularly, take it as soon as possible. If it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take double doses at once.

- *If you use too much (overdose)*

May cause abdominal distension. In the event of overdose, symptomatic treatment should be given.

While you are using it

- *Things you must do*

Follow your doctor's or pharmacist's instruction when you are taking HOMECARE BAROSCON DOUBLE ACTION SUSPENSION

- *Things you must not do*

Do not use this product after the expiry date which is stated on the label.

- *Things to be careful of*

Each 10ml (single sachet) dose has a sodium content of 127.9mg (5.56mmol) and 130mg (3.25mmol) of calcium. This should be taken into account when a highly restricted salt diet or calcium diet is recommended.

Side effects

Like all medicines, HOMECARE BAROSCON DOUBLE ACTION SUSPENSION can cause side effects, although not everybody gets them.

Very rarely (<1/10,000) patients sensitive to the ingredients may develop allergic manifestations such as urticaria or bronchospasm, anaphylactic or anaphylactoid reactions.

Ingestion of large quantities of calcium carbonate may cause alkalosis, hypercalcaemia, acid rebound, milk alkali syndrome or constipation. These usually occur following larger than recommended dosages.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers →Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of HOMECARE BAROSCON DOUBLE ACTION SUSPENSION

- *Storage*

Store at temperature below 30ºC, protect from sunlight & moisture.

- *Disposal*

HOMECARE BAROSCON DOUBLE ACTION should be disposed when your doctor or pharmacist tells you to stop taking this product or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required.

Product Description

- *What it looks like*

Peppermint-flavoured, pale yellow or cream-coloured suspensions

- *Ingredients*
  - Active ingredient(s)

Sodium Alginate
Calcium Carbonate
Sodium Bicarbonate

- Inactive Ingredients:

Methyl Parahydroxybenzoate
Propyl Parahydroxybenzoate

- *MAL number(s):*

MAL18046003X

Manufacturer

Yungjin Pharm. Co., Ltd.,
66, Muha-ro, Namyang-eup,
Hwaseong-si, Gyeonggi-do,
Republic of Korea.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3, 5, 7 & 12, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor,
Malaysia

Date of revision

08/08/2024

Serial Number

NPRA(R2/1)240808/1325