

XIGATIN FILM COATED TABLET 100MG

Sitagliptin phosphate monohydrate (eq. to sitagliptin 100mg)

What is in this leaflet

This leaflet contains information below:

1. What XIGATIN FILM COATED TABLET 100MG is used for
2. How XIGATIN FILM COATED TABLET 100MG works
3. Before you use XIGATIN FILM COATED TABLET 100MG
4. How to use XIGATIN FILM COATED TABLET 100MG
5. While you are using XIGATIN FILM COATED TABLET 100MG
6. Side effects
7. Storage and disposal of XIGATIN FILM COATED TABLET 100MG
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What XIGATIN FILM COATED TABLET 100MG is used for

XIGATIN FILM COATED TABLET 100MG is indicated as an addition to diet and exercise to improve blood glucose control in patients with type 2 diabetes mellitus.

How XIGATIN FILM COATED TABLET 100MG works

XIGATIN is a member of a class of medicines you take by mouth called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors) that lowers blood sugar levels in patients with type 2 diabetes mellitus. Type 2 diabetes is also called non-insulin-dependent diabetes mellitus, or NIDDM.

Before you use XIGATIN FILM COATED TABLET 100MG

When you must not use it

Do not take XIGATIN if you have allergy towards any of the components in XIGATIN.

Before you start to use it

Tell your doctor or pharmacist if you are allergic to any medicines or if have had:

- type 1 diabetes
- diabetic ketoacidosis (increased ketones in the blood or urine)
- any kidney problems, or any past or present medical problems
- an allergic reaction to XIGATIN

Taking other medicines

XIGATIN may be taken with most medicines. Tell your doctor about all the medicines you take. This includes prescription and non-prescription medicines, and herbal supplements.

How to use XIGATIN FILM COATED TABLET 100MG

How much to use

Take XIGATIN exactly as your doctor has prescribed. The recommended dose is to take:

- one 100 mg tablet
- once a day
- by mouth, with or without food

If you have kidney problems, your doctor may prescribe lower doses.

Your doctor may prescribe XIGATIN along with certain other medicines that lower blood sugar.

When to use it

May be taken with or without food. Take as directed by your doctor or pharmacist.

How long to use it

XIGATIN should be taken as long as your doctor advised you to do so that you can continue to help control your blood sugar.

Diet and exercise can help your body use its blood sugar better. It is important to stay on your doctor recommended diet, exercise and weight loss program while taking XIGATIN.

If you forgot to use it

If you miss a dose, take the missed dose as soon as you remember. However, if it's almost time for the next dose, skip the missed dose and continue the regular dosing schedule. Do not take a double dose to make up for a missed one.

If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using XIGATIN FILM COATED TABLET 100MG

Things you must do

Follow your doctor's instruction when you are taking XIGATIN.

Tell all the doctors, dentists and pharmacists treating you that you are taking XIGATIN.

You should promptly report to physician if you become pregnant while taking XIGATIN.

Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give this medication to anyone else, even if they have the same condition as you.

Things to be careful of

Cases of inflammation of the pancreas (pancreatitis) have been reported in patients receiving XIGATIN. Pancreatitis can be a serious, potentially life-threatening medical condition. Stop taking XIGATIN and call your doctor if you experience severe and persistent stomach pain, with or without vomiting, because you could have pancreatitis.

Cases of a skin reaction called bullous pemphigoid that can require treatment in a hospital have been reported in patients receiving XIGATIN. Tell your doctor if you develop blisters or the breakdown of your skin (erosion). Your doctor may tell you to stop taking XIGATIN.

Use in children

XIGATIN has not been studied in children under 18 years of age.

Use in pregnancy and breast-feeding

Women who are pregnant or plan to become pregnant should consult with their doctor before taking XIGATIN. XIGATIN is not recommended for use during pregnancy.

It is not known if XIGATIN passes into breast milk. You should not use XIGATIN if you are breast-feeding or plan to breast-feed.

Driving or using machines

There is no information to suggest that XIGATIN affects your ability to drive a car or operate machinery.

XIGATIN FILM COATED TABLET 100MG

Sitagliptin phosphate monohydrate (eq. to sitagliptin 100mg)

Side effects

Like all medicines, XIGATIN can cause side effects, although not everybody gets them. Tell your doctor or pharmacist if you notice anything else that making you feel unwell.

Additional side effects have been reported in general use with XIGATIN, by itself and/or with other diabetes medicines:

- Reduced number of platelets
- Allergic reactions, which may be serious, including rash, hives, and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing. If you have an allergic reaction, stop taking XIGATIN and call your doctor right away. Your doctor may prescribe a medication to treat your allergic reaction and a different medication for your diabetes.
- low blood sugar with symptoms (hypoglycemia)
- Headache
- Dizziness
- Upper respiratory infection
- Stuffy or runny nose and sore throat
- Constipation
- Vomiting
- Inflammation of the pancreas
- Itching
- Blisters
- Joint pain
- Muscle aches
- Back pain
- Arm or leg pain
- Kidney problems

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumer → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and disposal of XIGATIN FILM COATED TABLET 100MG

Storage

Store below 30°C, protect from light & moisture.


Disposal

Dispose when you stop taking the XIGATIN as instructed by your doctor or pharmacist or the tablets have passed their expiry date. Ask your pharmacist how to dispose medicines that are no longer required.

Product description

What it looks like

An orange brown color film coated tablet, one side impressed with the letter "100".



Ingredients

Active ingredient:-

Sitagliptin Phosphate Monohydrate

Inactive ingredients(s):-

Croscarmellose Sodium, Microcrystalline Cellulose, Anhydrous Dibasic Calcium Phosphate, PVP K-30, Sodium Stearyl Fumarate, Magnesium Stearate, Opadry II Brown

MAL number

MAL21086014AZ

Manufacturer and Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd. Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision

26/03/2026

Serial Number

NPRA(R2/1)250131/1413