

EZYCOR TABLET 10MG

Ezetimibe Tablet 10mg

What is in this leaflet

1. What Ezycor is used for
2. How Ezycor works
3. Before you use Ezycor
4. How to use Ezycor
5. While you are using Ezycor
6. Side effects
7. Storage and disposal of Ezycor
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Ezycor is used for

Ezycor is a tablet that contains 10mg of ezetimibe as the active ingredient.

- Ezycor is a cholesterol-lowering medicine.
- Ezycor is for patients whose cholesterol levels are too high and when diet alone cannot lower these levels adequately.
- Ezycor reduces elevated total-cholesterol, LDL (bad) cholesterol and triglycerides and increases HDL (good) cholesterol.
- Ezycor may be taken with other cholesterol-lowering medicines known as statins or alone, in addition to diet.
- Ezycor may be taken with fenofibrate, another cholesterol-lowering medicine, in addition to diet.
- For patients with heart disease, Ezycor combined with cholesterol-lowering medicines called statins reduces the risk of heart attack, stroke, surgery to increase heart blood flow, or hospitalization for chest pain.

If you have sitosterolemia, your doctor has prescribed Ezycor to lower the level of plant sterols in your blood.

Sitosterolemia is a hereditary illness that increases the levels of plant sterols in your blood.

How Ezycor works

- Ezycor works by decreasing the absorption of cholesterol in the small intestine. Statins and fenofibrate lower cholesterol in a different way; they work in the liver.
- Ezycor adds to the cholesterol-lowering effect of statins and fenofibrate.

Why has my doctor prescribed Ezycor?

Your doctor has prescribed Ezycor to reduce the amount of cholesterol and triglycerides in your blood. Cholesterol is one of several fatty substances found in the bloodstream. Your total cholesterol is made up mainly of LDL and HDL cholesterol.

LDL cholesterol is often called “bad” cholesterol because it can build up in the walls of your arteries forming plaque. Eventually this plaque build-up can lead to a narrowing of the arteries. This narrowing can slow or block blood flow to vital organs such as the heart and brain. This blocking of blood flow can result in a heart attack or stroke.

HDL cholesterol is often called “good” cholesterol because it helps keep the bad cholesterol from building up in the arteries and protects against heart disease.

Triglycerides are another form of fat in your blood that may increase your risk for heart disease.

How can high cholesterol be treated?

High cholesterol can be treated in two main ways:

Lifestyle Changes – including a cholesterol-lowering diet, increased physical activity, and weight management.

Medication – Cholesterol-lowering medicines are used together with lifestyle changes to help lower cholesterol. Your doctor has prescribed Ezycor to help lower your cholesterol.

Before you use Ezycor

It is important that you continue taking Ezycor daily as prescribed by your doctor.

Even if you are taking medication to treat high cholesterol, it is important to have your cholesterol measured regularly. You should also know your cholesterol levels and goals.

Use in elderly

No special precautions.

Use in children

Ezycor is not recommended in children under age 10.

- When you must not use it

Patients who are hypersensitive (allergic) to Ezycor or any of its ingredients should not take Ezycor.

- Before you start to use it

Tell your doctor about:

Pregnancy: If you are pregnant or plan to become pregnant, Ezycor may not be right for you.

Breast-feeding: If you are breast-feeding, Ezycor may be passed in your milk to your baby.

Medical Problems or Allergies:

Inform any medical problems (including liver disease or liver problems) or allergies you have now or have had in the past.

- Taking other medicines

You should always tell your doctor about all drugs that you are taking or plan to take, including those obtained without a prescription.

EZYCOR TABLET 10MG

Ezetimibe Tablet 10mg

In particular, tell your doctor if you are taking medicine(s) with any of the following active ingredients:

- Cyclosporin (often used in organ transplant patients)
- Medicines with an active ingredient to prevent blood clots, such as warfarin or fluindione (anticoagulants)
- Cholestyramine (also used to lower cholesterol), because it affects the way Ezycor works
- Fibrates (also used to lower cholesterol)

How to use Ezycor

- How much to use

Take one 10 mg tablet by mouth each day, at any time of day.

- When to use it

- Take Ezycor with or without food.
- Your doctor may let you take Ezycor along with other medicines, known as statins, or with another medicine known as fenofibrate, to help you better control your cholesterol.
- If you are taking a statin or fenofibrate, Ezycor can be taken at the same time as the other medicine.
- If your doctor has prescribed Ezycor along with cholestyramine (a bile acid sequestrant) or any other bile acid sequestrant, Ezycor should be taken at least 2 hours before or 4 hours after taking the bile acid sequestrant.

- How long to use it

Continue taking your cholesterol-lowering medicines unless your doctor tells you to stop.

- If you forget to use it

Try to take Ezycor as prescribed. However, if you miss a dose, just resume the usual schedule of one tablet daily.

- If you use too much (overdose)

Take Ezycor only as prescribed for you. If you take more Ezycor than you were prescribed, contact your doctor or pharmacist.

While you are using Ezycor

- Things you must do

- Ezycor should be taken as directed by your health care provider.
- Talk about any other medicines that you are taking or plan to take. Include those you can get without a prescription.
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not take Ezycor if you are hypersensitive (allergic) to Ezycor or any of its ingredients.
- Do not give Ezycor to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Can I drive or operate machinery while using Ezycor?

There have been side effects reported with Ezycor that may affect your ability to drive or operate machinery. Individual responses to Ezycor may vary. (See Side effects)

Side effects

Ezycor was generally well tolerated. The side effects were usually mild. Side effects generally do not cause patients to stop taking Ezycor.

When used alone, the following side effects were reported:

Common: abdominal pain, diarrhea, flatulence, feeling tired.

Uncommon: elevations in some laboratory blood tests of liver

(transaminases) or muscle (CK) function, cough, indigestion, heartburn, nausea, joint pain, muscle spasms, neck pain, decreased appetite, pain, chest pain, hot flush, high blood pressure.

Additionally, when used with a statin, the following side effects were reported:

Common: elevations in some laboratory blood tests of liver function (transaminases), headache, muscle pain, tenderness or weakness.

Uncommon: tingling sensation, dry mouth, itching, rash, hives, back pain, muscle weakness, pain in arms and legs, unusual tiredness or weakness, swelling, especially in the hands and feet.

When used with fenofibrate, the following side effect was reported:

Common: abdominal pain.

Additionally, the following side effects have been reported in general use: allergic reactions (which may require treatment right away) including swelling of the face, lips, tongue, and/or throat that may cause difficulty in breathing or swallowing, rash, and hives, raised red rash, sometimes with target-shaped lesions, muscle aches, unusual tiredness or weakness, alterations in some laboratory blood tests, liver problems, inflammation of the pancreas, constipation, dizziness, tingling sensation, depression, gallstones, and inflammation of the gallbladder.

Contact your doctor promptly if you experience unexplained muscle pain, tenderness, or weakness while taking Ezycor.

This is because on rare occasions, muscle problems can be serious,

EZYCOR TABLET 10MG

Ezetimibe Tablet 10mg

including muscle breakdown resulting in kidney damage.

Talk to your doctor anytime you have a medical problem you think may be related to Ezycor.

If you are prescribed Ezycor with a statin, your healthcare professional may want to conduct routine blood tests to check your liver function before and after starting treatment.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nppra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

How can I learn more about Ezycor and my condition?

You may obtain further information from your doctor or pharmacist.

Storage and Disposal of Ezycor

- Storage

Keep out of reach of children.

Store below 30°C. Protect from moisture.

- Disposal

Do not use this medicine after the date shown following expiry date on the container.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Product Description

- What it looks like

White to off white, capsule shaped, flat faced with beveled edge, uncoated tablets, debossed with “10” on one side and plain on other side.

- Ingredients

- Active ingredient
Ezetimibe

-Inactive ingredients

Lactose monohydrate
Croscarmellose sodium
Povidone k-30
Sodium lauryl sulfate
Polysorbate 80
Magnesium stearate

- MAL Number:

MAL XXXXXXAZ

Manufacturer

INTAS PHARMACEUTICALS LIMITED.

Plot No. 457 & 458, Village Matoda, Bavla Road and Plot No. 191/218P, Village: Chacharwadi, Tal- Sanand, Ahmedabad, Gujarat, INDIA.

Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5 & 7, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor Darul Ehsan, Malaysia.

Date of revision

10/10/2022

Serial Number

NPRA (R2) 22/129