

TORIXIB FILM COATED TABLET

Etoricoxib 60mg, 90mg, 120mg

What is in this leaflet

1. What Torixib Film Coated Tablet is used for
2. How Torixib Film Coated Tablet works
3. Before you use Torixib Film Coated Tablet
4. How to use Torixib Film Coated Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Torixib Film Coated Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision
11. Serial Number

What Torixib Film Coated Tablet is used for

Your doctor has prescribed Torixib Film Coated Tablet for treatment of:

- osteoarthritis
- rheumatoid arthritis
- ankylosing spondylitis
- chronic low back pain (60 mg only)
- acute pain after minor dental procedures
- menstrual pain
- gout attacks

Osteoarthritis is a joint disease. It results from the gradual breakdown of the cartilage that cushions the ends of the bones resulting in pain, inflammation, tenderness, stiffness, and physical disability.

Rheumatoid arthritis is a chronic disease of the joints that causes pain, stiffness, swelling and loss of function in the joints. It may also cause inflammation in other parts of the body organs.

Ankylosing spondylitis is an inflammatory disease of the spine and large joints.

Gout is a disorder characterised by sudden, recurring attacks of pain and inflammation in one or more joints.

You may obtain further information from your doctor or pharmacist, who have more detailed information.

How Torixib Film Coated Tablet works

Torixib Film Coated Tablet is a member of a group of medicines called selective COX-2 inhibitors that are used to reduce pain and inflammation. Torixib Film Coated Tablet is not a narcotic.

Before you use Torixib Film Coated Tablet

- When you must not use it

Do not use Torixib Film Coated Tablet if you:

- are allergic to etoricoxib or any other ingredients of Torixib Film Coated Tablet.
- have had heart failure, a heart attack, bypass surgery, chest pain, narrow or blocked arteries of the extremities, a stroke or mini stroke.
- have high blood pressure that has not been controlled by treatment.

- Before you start to use it

Tell your doctor about any medical problems or allergies you have now or have had, including:

- history of angina, heart attack or a blocked artery in your heart.
- narrow or blocked arteries of the extremities.
- kidney disease.
- liver disease.
- dehydration, for example by a prolonged bout of vomiting or diarrhea.
- have or have had ulceration, bleeding or perforation of your stomach or intestines.
- heart failure.
- high blood pressure.
- swelling due to fluid retention.
- an allergic reaction to aspirin or other anti-inflammatory medicines (commonly known as NSAIDs).

- a history of stroke or mini stroke.
- conditions which increase your risk of coronary artery disease or atherosclerosis such as high blood pressure, diabetes, high cholesterol or smoking.
- being treated for an infection. Torixib Film Coated Tablet can mask or hide a fever, which is a sign of infection.
- have had asthma, nasal inflammation, and/or itching or hives caused by Aspirin or NSAIDs.
- suffering from dyspepsia (indigestion).

Etoricoxib may cause severe skin reactions. Symptoms may include skin rash, blisters and peeling of the skin. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Use in children

Torixib Film Coated Tablet has not been adequately studied in children. Therefore, Torixib Film Coated Tablet should not be given to children.

Use in elderly

Torixib Film Coated Tablet works equally well in older and younger adult patients.

Adverse experiences may occur at a higher incidence in older patients compared to younger patients. If you are elderly (i.e., over 65 years of age), your doctor will want to appropriately keep a check on you. No dosage adjustment is necessary for older patients.

Pregnancy and breastfeeding

Tell your doctor if you are:

- attempting to become pregnant or become pregnant during treatment because Torixib Film Coated Tablet may cause fetal

TORIXIB FILM COATED TABLET

Etoricoxib 60mg, 90mg, 120mg

harm and should not be used during pregnancy.

- breastfeeding or plan to breastfeed.

- Taking other medicines

You should always tell your doctor about all medicines that you are taking or plan to take, including those obtained without a prescription.

Your doctor may want to check that your medicines are working properly if you are taking:

- warfarin (a blood thinner)
- rifampin (an antibiotic)
- diuretics (water tablets)
- ACE inhibitors and angiotensin receptor blockers (medicines used for high blood pressure and heart failure)
- lithium (a medicine used to treat a some type of depression)
- birth control pills
- hormone replacement therapy
- methotrexate (a medicine used for suppressing the immune system).

Torixib Film Coated Tablet can be taken with low-dose aspirin. If you are currently taking low-dose aspirin for prevention of heart attack or stroke, you should not stop taking aspirin until you talk to your doctor because Torixib Film Coated Tablet cannot substitute for aspirin for this purpose.

How to use Torixib Film Coated Tablet

- How much to use

Torixib Film Coated Tablet should be taken once a day. You may take Torixib Film Coated Tablet with or without food.

Your doctor will decide what dose of Torixib Film Coated Tablet you should take and how long you should take it.

Treatment of osteoarthritis:

- The recommended dose is 30mg or 60mg once a day.

Treatment of rheumatoid arthritis:

- The recommended dose is 60mg once a day, and may increase to 90mg once a day if needed.

Treatment of ankylosing spondylitis:

- The recommended dose is 60mg once a day, and may increase to 90mg once a day if needed.

Chronic low back pain:

- The recommended dose is 60 mg once daily.

For acute pain conditions, the recommended dose is 90mg or 120mg once daily. Torixib Film Coated Tablet should be used only for the acute period limited to a maximum of 8 days.

For the relief of gout attacks and treatment of menstrual pain:

- The recommended dose is 120mg once a day which should only be used for the acute painful period.

For the relief of pain after minor dental procedures:

- The recommended dose is 90mg once a day.

Doses greater than those recommended for each condition have either not demonstrated additional efficacy or have not been studied. Therefore, the daily doses stated above for each condition should not be exceeded.

If you have mild or moderate liver disease, your doctor might prescribe you a lower dose or recommend that you do not take Torixib Film Coated Tablet.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue using Torixib Film Coated Tablet for the duration that your doctor recommends.

- If you forget to use it

Try to take Torixib Film Coated Tablet as prescribed. If you miss a dose, do not take an extra dose. Just resume your usual schedule the following day.

- If you use too much (overdose)

If you take more than the prescribed dosage, contact your doctor immediately.

While you are using it

- Things you must do

Consult your doctor if:

- any of the following symptoms appear or worsen: shortness of breath, chest pains or ankle swelling. Stop your treatment with Torixib Film Coated Tablet and consult a doctor, as soon as is practical.
- you have kidney, liver or heart disease, your doctor will want to appropriately keep a check on you.
- you develop any symptoms that could indicate a severe allergic reaction such as inability to breathe or a serious skin reaction, you must consult a doctor on an urgent basis.

Your doctor will want to discuss your treatment from time to time. It is important that you use the lowest dose that controls your pain.

Torixib Film Coated Tablet can increase blood pressure in some people, especially in high doses, and this could increase the risk of heart attacks and strokes. Your doctor will want to check your blood pressure

TORIXIB FILM COATED TABLET

Etoricoxib 60mg, 90mg, 120mg

from time to time, to make sure that it is safe to continue treatment.

- *Things you must not do*

Do not share Torixib Film Coated Tablet with anyone else, it was prescribed only for you.

Do not take Torixib Film Coated Tablet for longer than necessary. This is because the risk of heart attacks and strokes might increase after prolonged treatment, especially with high doses.

- *Things to be careful of*

Driving or operating machinery

Be careful driving or operating machinery until you know how Torixib Film Coated Tablet affects you.

Side effects

Any medicine may have unintended or undesirable effects, so-called side effects. Like all medicines, Torixib Film Coated Tablet may cause side effects.

Some of the side effects reported in the studies included:

- dizziness,
- swelling of the legs and/or feet,
- weakness and fatigue,
- high blood pressure,
- nausea,
- heartburn,
- upset stomach,
- headache,
- dry socket (inflammation and pain after a tooth extraction).

Additionally, the following have been reported:

- allergic reactions (which may be serious enough to require immediate medical attention) including swelling of the face, lips, tongue, and/or throat which may cause difficulty in breathing or swallowing, wheezing, rash, itching and hives;
- redness of the skin,

- severe skin reactions which may occur without warning,
- taste alteration,
- abnormal rhythm of the heart,
- heart failure,
- palpitations,
- feeling of tightness, pressure or heaviness in the chest,
- stomach pain,
- stomach ulcers that may become serious and may bleed and may occur at any time during use and without warning,
- vomiting,
- liver problems including liver failure, yellowing of the skin and eyes (jaundice),
- serious kidney problems,
- high levels of potassium in your blood,
- insomnia,
- anxiety,
- depression,
- restlessness,
- drowsiness,
- mouth ulcers,
- diarrhea,
- severe increase in blood pressure,
- confusion,
- hallucinations,
- platelets decreased,
- blurred vision.

Other side effects may also occur rarely, and as with any prescription medicine, some side effects may be serious. Ask your doctor or pharmacist for more information. Both have a more complete list of side effects. Tell your doctor or pharmacist promptly about these or any other unusual symptoms.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Torixib Film Coated Tablet

- *Storage*

Store below 30°C. Store in the original package. Protect from light and moisture.

Keep Torixib Film Coated Tablet out of reach of children.

- *Disposal*

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- *What it looks like*

Torixib Film Coated Tablet 60mg: A dark green round film coated tablet.
 Torixib Film Coated Tablet 90mg: A white round film coated tablet, one side impressed with “Y”.
 Torixib Film Coated Tablet 120mg: A light green round film coated tablet.

- *Ingredients*

- Active ingredient:
Etoricoxib

- Inactive ingredients:

Core tablet

- Microcrystalline Cellulose
- Dicalcium Phosphate Anhydrous
- Croscarmellose Sodium
- Magnesium Stearate

Film coating

- Torixib Film Coated Tablet 60mg: Opadry II Green
- Torixib Film Coated Tablet 90mg: Opadry II White
- Torixib Film Coated Tablet 120mg: Opadry II Green

- *MAL number:*

MALXXXXXXXXXAZ

TORIXIB FILM COATED TABLET

Etoricoxib 60mg, 90mg, 120mg

Manufacturer and Product

Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7 & 12, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor,
Malaysia.

Date of revision

04/01/2024

Serial Number

NPRA (R2) 23/048