

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg

Hydroxychloroquine Sulphate (200 mg)

What is in this leaflet

1. What YSP
Hydroxychloroquine Sulphate Film Coated Tablet 200 mg is used for
2. How YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg works
3. Before you use YSP
Hydroxychloroquine Sulphate Film Coated Tablet 200 mg
4. How to use YSP
Hydroxychloroquine Sulphate Film Coated Tablet 200 mg
5. While you are using it
6. Side effects
7. Storage and Disposal of YSP
Hydroxychloroquine Sulphate Film Coated Tablet 200 mg
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision
11. Serial Number

What YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg is used for

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg can be used for:

- Rheumatoid arthritis (inflammation of the joints)
- Juvenile chronic arthritis (in children)
- Discoid and systemic lupus erythematosus (a disease of the skin or the internal organs)
- Skin problems which are due to/ aggravated by sunlight

How YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg works

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200mg contains a medicine called hydroxychloroquine sulphate. It works by reducing inflammation in people with autoimmune diseases (this is where the body's immune system attacks itself by mistake).

Before you use YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg

- When you must not use it

Do not take YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg if you:

- Are allergic to hydroxychloroquine, quinine or other 4-aminoquinoline compounds.
- Have signs of an allergic reaction include a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue.
- Have an eye problem which affects the retina, the inside of the eye (maculopathy) or you get a change in eye colour or any other eye problem
- Are pregnant, plan to get pregnant, breastfeeding or plan to start breastfeeding unless your doctor considers the benefits outweigh the risks.

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200mg may be associated with a small increased risk of malformations and should not be used during pregnancy unless your doctor considers the benefits outweigh the risks.

Do not use this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg.

- Before you start to use it

Take special care and check with your doctor if:

- You have liver or kidney problems.
- You have serious stomach or gut problems.
- You are taking a drug called Tamoxifen, used to treat breast cancer.

- You have any problems with your blood. You may have blood tests to check this.
- You have heart problems (signs include breathlessness and chest pain) which may require monitoring.
- You have any problems with your nervous system or brain.
- You have psoriasis (red scaly patches on the skin usually affecting the knees, elbows and scalp).
- You have had a bad reaction to quinine in the past.
- You have a genetic condition known as 'glucose-6-dehydrogenase deficiency'.
- You have a rare illness called 'porphyria' which affects your metabolism.
- You have or have had myasthenia (a disease with general muscle weakness including in some cases muscles used for breathing). You may notice aggravation of symptoms such as muscle weakness, difficulty in swallowing, double vision, drooping of the upper eyelid etc.
- Hydroxychloroquine may cause heart, kidney or muscle disorders. Please ask your doctor to inform you of signs and symptoms of drug induced phospholipidosis. Hydroxychloroquine may need to be stopped.
- Hydroxychloroquine can cause lowering of the blood glucose level. Please ask your doctor to inform you of signs and symptoms of low blood glucose levels. A check of the blood glucose level may be necessary.
- Hydroxychloroquine may cause heart rhythm disorders in some patients: caution should be taken when using hydroxychloroquine if you were born with or have family history of prolonged QT interval, if you have acquired QT prolongation (seen on ECG,

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg

Hydroxychloroquine Sulphate (200 mg)

electrical recording of the heart), if you have heart disorders or have a history of heart attack (myocardial infarction), if you have salt imbalance in the blood (especially low level of potassium or magnesium, see section "Taking other medicines"). If you experience palpitations or irregular heartbeat, you should inform your doctor immediately. The risk of heart problems may increase with increase of the dose. Therefore, the recommended dosage should be followed.

- Serious skin rashes have been reported with the use of hydroxychloroquine. Frequently, the rash can involve ulcers of the mouth, throat, nose, genitals and conjunctivitis (red and swollen eyes). These serious skin rashes are often preceded by flu-like symptoms such as fever, headache and body ache. The rash may progress to widespread blistering and peeling of the skin. If you develop these skin symptoms, stop taking hydroxychloroquine and contact your doctor immediately.
- You have lactose intolerance.
- Before you take YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg you should have your eyes examined. This testing should be repeated at least every 12 months whilst taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg.
- If you are over 65, need to take a high dose (2 tablets a day) or have kidney problems, then this examination should be performed more often.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg.

- Taking other medicines

Tell your doctor, if you are taking the following medicines which may increase the chance of you getting side effects:

- Some antibiotics used for infections
- Cimetidine - used for stomach ulcers
- Neostigmine and pyridostigmine – used for muscle weakness (myasthenia gravis)
- Medicines that may affect the kidney or liver
- Medicines that affect the skin or the eyes
- Medicines known to affect the rhythm of your heart. This includes medicines used for abnormal heart rhythm (antiarrhythmics), for depression (tricyclic antidepressants), for psychiatric disorders (antipsychotics), for bacterial infections, or against malaria (e.g., halofantrine)
- Medicines for lipid disorders (such as gemfibrozil, statins)
- Medicines for depression (fluoxetine, paroxetine)

The following medicines can change the way YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg works or YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg may affect the way some of these medicines work:

- Digoxin - used for heart problems
- Medicines for epilepsy
- Medicines for diabetes (such as insulin/ oral antidiabetic drugs)
- Antacids - used for heartburn or indigestion. You should leave a gap of at least 2 hours between taking these medicines and YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg.
- Rabies vaccine
- Praziquantel
- Ciclosporin

- Agalsidase
- Itraconazole
- Clarithromycin
- Rifampicin
- Ritonavir
- Grapefruit juice

How to use YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg

Always take YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

- Swallow the tablets orally.
- Take this medicine with a meal or a glass of milk.

- How much to use

Adults (including the elderly): 1-2 tablets per day.

- The dose can be reduced from 400 mg daily to 200 mg when there is no further improvement.
- Children: 1 tablet per day.
- This medicine is only suitable for children whose ideal body weight is more than 31 kg.

It may take several weeks before you notice the benefit of taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg, however, side effects may occur relatively early.

- When to use it

Take YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg at about the same time each day unless your doctor tells you otherwise. Taking your medicine at the same time each day will have the best effect. It will also help you remember when to take the medicine. If you are not sure when to use it, ask your doctor or pharmacist.

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg

Hydroxychloroquine Sulphate (200 mg)

- How long to use it

Continue taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg as long as your doctor recommends. You should not stop taking this medicine without medical advice.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you stop using it

Keep taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg until your doctor tells you to stop. Do not stop taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg just because you feel better. If you stop, your illness may get worse again.

- If you use too much (overdose)

If you take more YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg than you should, tell a doctor or go to a hospital casualty department straight away. Take the medicine pack with you. The following effects may happen: headache, problems with your eyesight, low potassium, fall in blood pressure, convulsions (fits), heart problems, leading to uneven heart beats, followed by sudden severe breathing problems and possibly heart attack. Overdose of YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg may have a fatal outcome. Young children and babies are particularly at risk if they accidentally take YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg. Take the child to a hospital straight away.

While you are using it

- Things you must do

Tell your doctor immediately and

- Stop taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg if you are pregnant, might become pregnant or think you may be pregnant.

- Stop taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg if you are breast-feeding or planning to breastfeed.

- Stop taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg immediately if any disturbances of vision (including abnormal colour vision).

- Things you must not do

- Do not give this medicine to anyone else, even if they have the same condition as you.
- Do not use this medicine to treat any other complaints unless your doctor tells you to.
- Do not stop taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg or change the dosage, without checking with your doctor.
- Avoid taking grapefruit juice as it may increase the risk of side effects.

- Things to be careful of

Driving and using machines

You may get eye problems while taking this medicine. If you experience these side effects, do not drive or use any tools or machines, and tell your doctor straight away.

Blood tests

Periodic blood counts are advised for patients on long-term therapy and YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg should be discontinued if abnormalities occur.

A blood test may show changes in the way the liver is working and occasionally the liver may stop working.

Some people being treated with YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg can experience mental health problems such as irrational thoughts, anxiety, hallucinations, feeling confused or feeling depressed, including thoughts of self-harm or suicide, even those who have never had similar problems before. If you or others around you notice any of these side effects seek medical advice straight away.

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg contains lactose. If you have been told by your doctor that you cannot tolerate some sugars, talk to your doctor before taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg.

Side effects

All medicines can cause side effects although not everybody gets them. Do not be alarmed by this list of possible side effects. You may not experience any of them.

Stop taking YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg and see a doctor straight away if you notice any of the following serious side effects:

Eye problems

- This includes changes in the colour of your eye and problems with your eyesight such as blurring, sensitivity to light or the way you see colour.

Skin effects

- Rash with a fever and flu-like symptoms and enlarged lymph nodes. This could be a condition called Drug Reaction with

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg

Hydroxychloroquine Sulphate (200 mg)

Eosinophilia and Systemic Symptoms (DRESS).

- Blistering, widespread scaly skin, pus-filled spots together with fever. This could be a condition called Acute Generalized Exanthematous Pustulosis (AGEP).
- Blistering or peeling of the skin around the lips, eyes, mouth, nose and genitals, flu-like symptoms and fever. This known as Stevens-Johnson syndrome (SJS).
- Multiple skin lesions, itching of the skin, joint aches, fever and a general ill feeling. This could be a condition called Toxic Epidermal Necrolysis (TEN).
- Skin reaction including plum-coloured, raised, painful sores, particularly on your arms, hands, fingers, face and neck, which may also be accompanied by fever. This could be a condition called Sweet's syndrome.
- Skin rashes, itching, changes in the colour of your skin or the inside of your nose or mouth.
- Psoriasis (red scaly patches on the skin usually affecting the knees, elbows and scalp).
- Hair loss or loss of hair colour.
- You have an allergic reaction. The signs may include: a red or lumpy rash, swallowing or breathing problems, swelling of your eyelids, lips, face, throat or tongue.

Stomach and intestinal problems

- Diarrhoea, nausea, loss of appetite (anorexia) or stomach pain.

Brain problems

- Feeling nervous, ringing in the ear (tinnitus), hearing loss, headache, fits, balance problems (vertigo) or feeling dizzy.
- Feeling depressed, feeling nervous or anxious, feeling confused, agitated, difficulty

sleeping, delusions, hallucinations, changes in mood, feeling elated or overexcited. Not known (frequency cannot be estimated from the available data).

- Common: mood swings

Muscle problems

- You have any muscle weakness, cramps, stiffness or spasms or changes in sensation such as tingling.
- Trembling, muscle spasm or irregular jerky movements.

Blood problems

- You may get infections more easily than usual. This could be due to bone marrow depression or a blood disorder called 'agranulocytosis'.
- You may bruise more easily than usual. This could be due to a blood problem called 'thrombocytopenia'.
- You feel tired, faint or dizzy and have pale skin. These could be 'anaemia'.
- You feel weak, short of breath, bruise more easily than usual and get infections more easily than usual. These could be 'aplastic anaemia'.

Heart problems

- Weakening of the heart muscle (cardiomyopathy) resulting in difficulty in breathing, coughing, high blood pressure, swelling, increased heart rate, low amount of urine.
- Abnormal heart rhythm, life-threatening irregular heart rhythm (seen on ECG) (frequency cannot be estimated from the available data).

Liver problems

- Liver problems that may cause the eyes or skin to go yellow (jaundice).

Others

- Having thoughts of self-harm or suicide (see *Before you start to use it*).
- Symptoms of a condition called 'porphyria' which may include stomach pain, being sick, fits, blisters, itching.
- Symptoms of lowering of the blood glucose level (hypoglycaemia). You may feel a sense of nervousness, shaky or sweaty.
- Accumulation of a type of fat in tissue causing harm. The doctor may decide to stop the treatment with YSP Hydroxychloroquine Sulphate Film Coated Tablet 200mg.

Talk to your doctor or pharmacist if you get any of the side effects or if you notice any side effects not listed in this leaflet.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of YSP Hydroxychloroquine Sulphate Film Coated Tablet 200mg

If you have any queries about any aspect of your medicine, or any questions regarding the information in this leaflet, discuss them with your doctor or pharmacist.

- Storage

Keep out of reach and sight of children. Store at temperature below 30°C. Protect from light and moisture.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to

YSP Hydroxychloroquine Sulphate Film Coated Tablet 200 mg


Hydroxychloroquine Sulphate (200 mg)

dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

A white round film coated tablet, one side impressed with “Y” and plain on the other side.



- Ingredients

Active ingredient:

Hydroxychloroquine sulphate

Inactive ingredients:

Lactose

Corn starch

Polyvinylpyrrolidone

Magnesium stearate

Hydroxypropyl methylcellulose

Polyethylene glycol

Dimethylpolysiloxane

Titanium dioxide

- MAL number

MAL25036165AZ

Manufacturer and Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
(199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7,
Section 13, Kawasan Perindustrian
Bandar Baru Bangi, 43000 Kajang,
Selangor, Malaysia.

Date of revision

05/09/2025

Serial Number

NPRA (R2) 24/045