

Barica Hard Capsule

Pregabalin 75 mg & 150 mg

What is in this leaflet

1. What Barica is used for
2. How Barica works
3. Before you use Barica
4. How to use Barica
5. While you are using it
6. Side effects
7. Storage and disposal of Barica
8. Product description
9. Manufacturer and Product registration holder
10. Date of revision
11. Serial number

What Barica is used for

Barica is used to treat:

• Neuropathic pain

Pregabalin is indicated for the treatment of peripheral and central neuropathic pain in adults.

• Epilepsy

Pregabalin is indicated as adjunctive therapy in adults with partial seizures with or without secondary generalisation.

• Generalised anxiety disorder

Pregabalin is indicated for the treatment of Generalised Anxiety Disorder (GAD) in adults.

• Fibromyalgia

Pregabalin is indicated for the management of fibromyalgia.

How Barica works

Barica helps decrease the number of nerve signals, and as a result calms down overly sensitive nerve cell.

Before you use Barica

- When you must not use it

Do not start taking Barica if you are allergic to pregabalin or any of the ingredients in Barica. (listed in section "Product description").

Symptoms of an allergic reaction to Barica may include:

- Serious, even life-threatening allergic reactions
- Suicidal thoughts or actions
- Swelling of your hands, legs and feet
- Dizziness and sleepiness

Pregnancy and breast-feeding

Animal studies have shown that pregabalin, the active ingredient in Barica, causes reproductive toxicity. Hence it should not be taken during pregnancy. Effective contraception must be used in women of child bearing potential.

Barica is excreted in the milk of lactating women. As the safety of Barica in infants is not known, breast-feeding is not recommended during treatment with Barica. You and your healthcare provider should discuss whether you should take Barica or breast-feed, but you should not do both.

- Before you start to use it

Tell your healthcare provider about all your medical conditions including if you:

- have or had depression, mood problems or suicidal thoughts or behavior
- have kidney problems or get kidney dialysis
- have heart problems including heart failure
- have a bleeding problem or a low blood platelet count
- have abused prescription medicines, or street drugs in the past
- have ever had swelling in your face, mouth (tongue, lips, gums) and neck (throat and larynx; angioedema)
- are in the first trimester of pregnancy as Barica may cause major birth defects in the unborn child
- have or had indigestion to milk or milk products (lactose intolerance)
- have brain malfunction (encephalopathy)
- have had cases of misuse, abuse and dependence (drug-seeking behavior) in the past
- are undergoing treatment of central neuropathic pain due to spinal cord injury
- have existing slow and ineffective breathing, or if you are taking a group of medicines called CNS depressants, including opioids

- Taking other medicines

Tell your healthcare provider about all the medicines you take, including prescription

and non-prescription medicines, vitamins or herbal supplements. Barica and other medicines may affect each other causing side effects.

Especially tell your healthcare provider if you take:

- Angiotensin converting enzyme (ACE) inhibitors, which are used to treat many conditions, including high blood pressure. You may have a higher chance for swelling and hives if these medicines are taken with Barica. See Section 5 "While you are using it".
- Avandia (rosiglitazone), Avandamet (contains rosiglitazone and metformin), or Actos (pioglitazone) for diabetes. You may have a higher chance of weight gain or swelling of your hands or feet if these medicines are taken with Barica. See Section 6 "Side effects".
- Any medicines that make you sleepy.
- Barica may increase the effects of ethanol and lorazepam.
- Barica appears to increase certain effects (impairment of thought process and gross bodily movements) caused by oxycodone.
- There are reports of lung failure, coma and deaths in patients taking Barica and other central nervous system depressant medications, including in patients who are drug abusers.

Know the medicines you take. Keep a list of them with you to show your healthcare provider and pharmacist each time you get a new medicine. Do not start a new medicine without talking to your healthcare provider.

How to use Barica

- How much to use

- Take Barica exactly as prescribed. Your healthcare provider will tell you how much Barica to take.
- Barica may be taken with or without food.
- Your healthcare provider may change your dose. Do not change your dose without talking to your healthcare provider.

Barica Hard Capsule

Pregabalin 75 mg & 150 mg

- The dose range is 150 to 600 mg per day given in either two or three divided doses.

- When to use it

Take Barica exactly as prescribed. Your healthcare provider will tell you when to take Barica. It is advised to take Barica at the same time each day.

- How long to use it

Do not stop taking Barica without checking with your healthcare provider. If you stop taking Barica suddenly, you may have trouble sleeping, nausea, headaches, anxiety, excessive sweating, and diarrhea. If you have epilepsy and you stop taking Barica suddenly, you may have seizures more often. If Barica is discontinued, taper the drug gradually over a minimum of 1 week rather than discontinue the drug abruptly. Talk with your healthcare provider about how to stop Barica slowly.

- If you forget to use it

If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, just skip the missed dose. Take the next dose at your regular time. Do not take two doses at the same time.

- If you use too much (overdose)

If you take too much Barica, call your healthcare provider or poison control center, or go to the nearest emergency room right away.

These side effects were reported when Barica was taken too much:

Affective disorder (abnormal disturbances of mood), sleepiness, confusion, depression, agitation and restlessness.

Fits (seizures) were also reported when the medicine was taken in overdose.

While you are using it

- Things you must do

Stop taking Barica and call your healthcare provider immediately if you have any of these signs of a serious allergic reaction:

- Swelling of your face, mouth, lips, gums, tongue, throat or neck
- Trouble breathing

- Rash, hives (raised bumps) or blisters

Call your healthcare provider immediately if you have any of these symptoms, especially if they are new, worse or worry you:

- Thoughts about suicide or dying
- Attempts to commit suicide
- New or worse depression
- New or worse anxiety
- Feeling agitated or restless
- Panic attacks
- Trouble sleeping (insomnia)
- New or worse irritability
- Acting aggressive, being angry or violent
- Acting on dangerous impulses
- An extreme increase in activity and talking (mania)
- Other unusual changes in behaviour or mood

- Things you must not do

Do not drive a car, work with machines, or do other dangerous activities until you know how Barica affects you.

Pregabalin may cause dizziness and extreme sleepiness which may impair the ability to perform tasks such as driving or operating machinery and could increase the occurrence of accidental injury (fall) in the elderly population.

Ask your healthcare provider about when it will be okay to do these activities.

- Things to be careful of

If you have suicidal thoughts or actions, do not stop Barica without first talking to a healthcare provider.

Stopping Barica suddenly can cause serious problems.

Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

How can I watch for early symptoms of suicidal thoughts and actions?

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.
- Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

Side effects

Very Common:

- Dizziness, sleepiness
- Headache

Common:

- Cold
- Increased appetite
- Elevated mood, confused state, irritability, depression, loss in sense of direction, inability to sleep, reduced sex drive
- Lack of voluntary coordination of muscle, loss in control of posture, muscle contraction, loss of memory, memory impairment, disturbance in attention, abnormal tingling and pricking sensation, reduced sense of touch, relaxed state/reduction of irritability or agitation, balance disorder, lack of energy
- Blurry vision and double vision
- Loss of balance
- Vomiting, constipation or hard bowels, accumulation of gas in alimentary canal, accumulation of gas or air in the abdomen, dry mouth
- Muscle cramp, joint pain, back pain, pain in the limbs, tightening of muscles
- Accumulation of fluid in tissues, abnormal walking, fall, feeling drunk, feeling abnormal, weakness
- Increased weight
- Diarrhea, vomiting

Rare:

- Stevens-Johnson syndrome

If you have diabetes, you should pay attention to your skin while taking Barica and tell your healthcare provider about any sores or skin problems.

Barica Hard Capsule

Pregabalin 75 mg & 150 mg

Tell your healthcare provider about any side effect that bothers you or that does not go away.

These are not all the possible side effects of Barica. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction

Monitoring by visiting the website npra.gov.my [Consumer → Reporting → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and disposal of Barica

- Storage

Store below 30°C.

Keep in a cool, dry place away from sunlight.

- Disposal

Safely throw away any Barica that is out of date or no longer needed.

Keep Barica and all medicines out of the reach of children.

Product Description

- What it looks like

Barica Hard Capsule 75mg: A size #4 red color cap and white color body hard gelatin capsule, cap and body imprinted with “YSP” on one side and “75” on another side, containing white powder.

Barica Hard Capsule 150mg: A size #2 white color hard gelatin capsule, cap and body imprinted with “150” on one side and “YSP” on the other side, containing white powder.

- Ingredients

- **Active ingredient**

Pregabalin

- **Inactive ingredients**

Lactose, pregelatinized starch and talc

MAL number:

MAL

Manufacturer and Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision

28/07/2025

Serial Number

NPRA (R2) 25/055