

CERITEC SYRUP 1MG/ML

Cetirizine Hydrochloride 1mg / ml

What is in this leaflet

This leaflet answers some common questions about **CERITEC SYRUP**. It does not contain all the available information. Please talk to healthcare professionals for further information.

What CERITEC syrup is used for

Symptomatic treatment of seasonal rhinitis, perennial allergic rhinitis and urticaria of allergic origin in adults and children ≥ 1 year.

How CERITEC Syrup works

Cetirizine inhibits allergic response, reduces bronchial hyperreactivity to histamine in the asthmatic patients and allergic reaction induced by specific allergens.

Before you use CERITEC Syrup

When you must not take it

You must not take it if you are having hypersensitivity towards Cetirizine, if you have severe renal impairment or you are breastfeeding.

Before you start to take it

Tell your doctor if you are taking any other medicine.

Taking other medicines

There are no known interactions with other drugs. Nevertheless, this product should be used with caution if sedatives are also being taken.

How to take CERITEC Syrup

How much to take

Please consult your doctor for more information on how much medicine to take.

When to take it

CERITEC Syrup is to be taken with or without food preferably in the evening.

How long to take it

CERITEC syrup should be taken according to doctor's prescription.

If you forgot to take it

If you miss a dose of CERITEC Syrup and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is nearing time for the next dose, do not double the dose to catch up, unless advised by your health care provider. Do not take 2 doses at once.

If you take too much

Drowsiness can be a symptom. Quickly seek for medical attention if you think that you or anyone else have taken too much CERITEC Syrup.

While you are using CERITEC Syrup

Things you must do

Take this medicine according to doctor's prescription

Things you must not do

Caution should be exercised when driving a car, operating potentially dangerous machinery, and if alcohol is taken concomitantly with CERITEC syrup.

Things to be careful of

CERITEC Syrup should not be administered to pregnant women during the first 3 months of pregnancy. This product is contraindicated in lactating women since cetirizine is excreted in breast milk. Use with caution in epileptic patients and patients at risk of convulsions. This product contains aspartame and is unsuitable for phenylketourics.

Side effects

Headache, dizziness, drowsiness, agitation, dry mouth, gastrointestinal discomfort and symptoms of hypersensitivity occurs occasionally.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects of adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

After using CERITEC syrup

Storage

Store below 30°C. Protect from light and moisture.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

What it looks like

Clear and colorless solution with banana flavor.

Ingredients

Cetirizine Hydrochloride.
Methyl Paraben and Propyl Paraben as preservatives

MAL number:

MAL12055023AZ

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor , Malaysia.

Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor , Malaysia.

Date of revision

01/08/2025

Serial Number

NPRA(R2/1)250731/1515