SALBUTAMOL-G EXPECTORANT

Salbutamol sulfate (1.2mg/5ml) , Guaifenesin (50mg/5ml)

What is in this leaflet

1. What SALBUTAMOL-G EXPECTORANT is used for
2. How SALBUTAMOL-G EXPECTORANT works
3. Before you use SALBUTAMOL-G EXPECTORANT
4. How to use SALBUTAMOL-G EXPECTORANT
5. While you are using it
6. Side effects
7. Storage and disposal of SALBUTAMOL-G EXPECTORANT
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What SALBUTAMOL-G EXPECTORANT is used for

Help to reduce thickness of phlegm for ease to cough up. To opening up the airways tubes in your lungs when your chest is tight or when you are wheezing.

How SALBUTAMOL-G EXPECTORANT works

Salbutamol is a bronchodilator which helps to open up the airways and relieve chest tightness, wheezing and cough in asthmatic patient. Guaifenesin is an expectorant which facilitates the removal of respiratory tract secretion. The combination of salbutamol with Guaifenesin is designed to relieve respiratory obstruction and ease breathing difficulties.

Before you use SALBUTAMOL-G EXPECTORANT

- *When you must not use it*

You must not take SALBUTAMOL-G EXPECTORANT if you are allergic towards salbutamol sulfate and/or Guaifenesin.

- *Before you start to use it*

Consult your doctor and pharmacist regarding the consumption of SALBUTAMOL-G EXPECTORANT if you are pregnant or breastfeeding. Tell your doctor or pharmacist if you are taking any other medicines or you have any other health related problems.

- *Taking other medicines*

Concurrent use of SALBUTAMOL-G EXPECTORANT with propranolol (blood pressure lowering medication) and MAOIs (MonoAmine Oxidase Inhibitors) (antidepressants) may cause unwanted effects.

Seek for your doctor or pharmacist advice if you are taking any of the medication mentioned or other medicines, including those you have bought without a prescription.

How to use SALBUTAMOL-G EXPECTORANT

- *How much to use*

Adults: Oral, 10 to 20 ml two to three times a day.
Children: 2 to 6 years: Oral, 5 to 10 ml two times a day.
Children: 7 to 12 years: Oral, 10ml two times a day.
Children: >12 years: Oral, 10ml to 20ml two times a day.

- *When to use it*

SALBUTAMOL-G EXPECTORANT should be taken according to the time advised by your doctor or pharmacist. It should be taken orally on an empty stomach which is 1 hour before or 2 hours after meals.

- *How long to use it*

SALBUTAMOL-G EXPECTORANT should be taken regularly and as long as your doctor advised you to do so.

- *If you forget to use it*

If you miss a dose of SALBUTAMOL-G EXPECTORANT and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

- *If you use too much (overdose)*

Quickly seek for medical attention immediately if you think that you or anyone else has taken too much SALBUTAMOL-G EXPECTORANT.

While you are using it

- *Things you must do*

Follow your doctor's instruction when you are taking SALBUTAMOL-G EXPECTORANT.

- *Things you must not do*

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give SALBUTAMOL-G EXPECTORANT to anyone else, even if they have the same condition as you.

- *Things to be careful of*

Increasing use of SALBUTAMOL-G EXPECTORANT may indicate deterioration of asthma control. Under this condition, please consult your doctor for further treatment plan. It should be administered cautiously to patients with thyroid disorders and diabetes. SALBUTAMOL-G EXPECTORANT should not be the only or main treatment in patient with severe or unstable asthma.

Side effects

SALBUTAMOL-G EXPECTORANT may cause headache, nausea, shaky or tense feeling, heart beats faster than usual.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of SALBUTAMOL-G EXPECTORANT

- *Storage*

Store below 30ºC. Protect from light and moisture.

- *Disposal*

SALBUTAMOL-G EXPECTORANT should be disposed of when your doctor or pharmacist tells you to stop taking this solution or it has passed the expiry date. It should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines that are no longer required. These measures will help to protect the environment.

Product description

- *What it looks like*

SALBUTAMOL-G EXPECTORANT is a clear and colorless solution with blackcurrant flavor.

- *Ingredients*
  - Active ingredient
    Salbutamol sulfate
    Guaifenesin.

- Inactive ingredients
  Sucralose
  Sodium Saccharin
  Sodium Benzoate
  Sodium Citrate
  Citric Acid
  Blackcurrant flavour

- *MAL number:*
  MAL14115013AZ

Manufacturer and Product Registration Holder
Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Date of revision
12/06/2025