

EDYFIL FILM COATED TABLET

Sildenafil (100mg, 50mg)

What is in this leaflet

1. What EDYFIL FILM COATED TABLET is used for
2. How EDYFIL FILM COATED TABLET works.
3. Before you use EDYFIL FILM COATED TABLET.
4. How to take EDYFIL FILM COATED TABLET.
5. While you are using EDYFIL FILM COATED TABLET.
6. Side effects.
7. Storage and disposal of EDYFIL FILM COATED TABLET
8. Product description.
9. Manufacturer and Product Registration Holder.
10. Date of revision

What EDYFIL FILM COATED TABLET is used for

EDYFIL FILM COATED TABLET treat erectile dysfunction, which is the inability to achieve or maintain a penile erection sufficient for satisfactory sexual performance.

How EDYFIL FILM COATED TABLET works

Sildenafil enhances the effect of Nitric oxide (NO) by inhibiting the phosphodiesterase type 5 (PDE5), which is responsible for degradation of cGMP in the corpus cavernosum.

The increased levels of cGMP in the corpus cavernosum, resulting in smooth muscle relaxation and inflow of blood to the corpus cavernosum and caused to erection of the penis

Before you use EDYFIL FILM COATED TABLET

-When you must not use it

When you are allergic to EDYFIL FILM COATED TABLET or any other ingredient listed at the end of this leaflet. Please avoid taking this medication if you are concurrently using nitric oxide donors, organic nitrates or organic nitrites in any form either regularly or intermittently

-Before you start to use it

Tell your doctor if you are allergic to any other medicines or have any other medical conditions.

-Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines may be affected by EDYFIL FILM COATED TABLET, or may affect how well it works:

- Cimetidine (used to treat ulcers).
- Some medicines used to treat fungal infections including ketoconazole or itraconazole.
- Some medicines used to treat HIV including ritonavir.
- Doxazosin or medicines called alpha blockers, which are used in the treatment of high blood pressure or prostate problems.
- Amlodipine (used to treat high blood pressure).

How to take EDYFIL FILM COATED TABLET

-How much to use

The recommended dose is 50 mg taken, as needed, approximately 1 hr before sexual activity. The maximum recommended daily dose is 100 mg. The maximum recommended dosing frequency is once per day. Please take this medication according to doctor's instructions.

-When to use it

EDYFIL FILM COATED TABLET should be taken according to doctor's instruction.

-How long to use it

EDYFIL FILM COATED TABLET should be taken according to doctor's prescription.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it. You should not take EDYFIL FILM COATED TABLET more than once a day.

-If you take too much (overdose)

Quickly seek for medical attention if you think that you or anyone else has taken too much EDYFIL FILM COATED TABLET.

While you are using EDYFIL FILM COATED TABLET

-Things you must do

Take the tablet according to doctor's prescription.

-Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction of your doctor. Do not give EDYFIL FILM COATED TABLET to anyone else, even if they have the same condition as you.

-Things to be careful of

Be careful drinking alcohol while taking EDYFIL FILM COATED TABLET. Alcohol can temporarily affect the ability to get an erection. Avoid drinking large quantities of alcohol before sexual activity.

If you experience dizziness, feeling tired or sick when taking EDYFIL FILM COATED TABLET, do not drive or operate machinery and contact your doctor immediately.

EDYFIL FILM COATED TABLET offer no protection against sexually transmitted diseases like HIV infection. Protective measures to be taken as advised by your doctor.

Side effects

The adverse events were generally transient and mild to moderate in nature. The incidence of some adverse events increased with dose. The most commonly reported adverse reactions were headache and flushing. Other commonly reported side effects are dizziness, blurred vision or sensitive towards light, transient abnormal coloured vision, palpitation, rhinitis and indigestion. Possible side effects includes hypersensitivity reaction such as skin rash, tachycardia (abnormal rapid heart rate), hypotension (low blood pressure), syncope (loss consciousness), epistaxis (nose bleeding), vomiting, eye pain, red eyes/bloodshot eyes, prolonged and painful erection.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of EDYFIL FILM COATED TABLET**Serial number**

NPRA(R2/1)240314/1225

-Storage

Keep in a tight container. Store below 30°C, protect from light & moisture.

-Disposal


Ask your pharmacist how to dispose medicine that no longer required or has passed expiry date. These measures will help to protect the environment.

Product description**-What it looks like****Edyfil Film Coated Tablet 100mg:**

A blue film coated elliptical tablet, one side impressed with 'E10' and the other side impressed with a score.

**Edyfil Film Coated Tablet 50mg:**

A blue film coated elliptical tablet, one side impressed with 'E5' and the other side impressed with a score.

**Ingredients****-Active Ingredient:**

Sildenafil Citrate

-Inactive Ingredients:

Microcrystalline Cellulose

Croscarmellose Sodium

Dicalcium Phosphate

Magnesium Stearate

Hydroxypropyl Cellulose

Opadry Blue

-MAL number(s)

Edyfil Film Coated Tablet 100mg -
MAL16125030AZ

Edyfil Film Coated Tablet 50mg -
MAL16125031AZ

Manufacturer

Y.S.P Industries (M) Sdn. Bhd.

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section
13, Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor,
Malaysia.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section
13, Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor,
Malaysia.

Date of revision

28/07/2026