

VALFIX FILM COATED TABLET 100MG

Lamivudine (100mg)

What is in this leaflet

1. What VALFIX F.C. TABLET is used for
2. How VALFIX F.C. TABLET works
3. Before you use VALFIX F.C. TABLET.
4. How to take VALFIX F.C. TABLET.
5. While you are using VALFIX F.C. TABLET
6. Side effects
7. Storage and disposal of VALFIX F.C. TABLET
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What VALFIX F.C. TABLET is used for
Treatment of patients \geq 16 years of age with chronic hepatitis B and evidence of hepatitis B virus (HBV) replication.

How VALFIX F.C. TABLET works

VALFIX F.C. TABLET is an antiviral medicine that suppresses the hepatitis B virus and belongs to a group of medicines called nucleoside analogue reverse transcriptase inhibitors (NRTIs).

Before you use VALFIX F.C. TABLET

When you must not take it

When you are allergic to VALFIX F.C. TABLET or any other ingredient listed at the end of this leaflet. Check with your doctor if you think this applies to you.

Before you start to take it

Tell your doctor if you are allergic to any other medicines or have any other medical conditions.

If you have Kidney Disease:

Discuss the use of VALFIX F.C. TABLET with your doctor or health care provider if you have kidney disease.

If you have Liver Disease/Liver Transplantation:

Please speak with your doctor or health care provider if you have a history of liver disease. In the case where treatment with VALFIX F.C. TABLET is discontinued for any reason, patient should periodically monitor for liver function.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from pharmacy, supermarket or health food shop.

Some medicines when taken with VALFIX F.C. TABLET may interfere with each other. These include:

- Zalcitabine and Emtricitabine, medicine to treat HIV infection.
- Other medicinal products containing lamivudine.

How to take VALFIX F.C. TABLET

How much to take

100mg to be taken once daily with or without food. Please take this medication according to doctor's instructions.

When to take it

VALFIX F.C. TABLET should be taken according to doctor's instruction.

How long to take it

VALFIX F.C. TABLET should be taken according to doctor's prescription.

If you forget to take it

If you miss a dose of VALFIX F.C. TABLET and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is nearing time for the next dose, do not double the dose to catch up, unless advised by your health care provider. Do not take 2 doses at once.

If you take too much (overdose)

Accidentally taking too much VALFIX F.C. TABLET is unlikely to cause any serious problems. However, you should seek for medical attention.

While you are using VALFIX F.C. TABLET

Things you must do

Take the tablet according to doctor's prescription.

Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction of your doctor. Do not give VALFIX F.C. TABLET to anyone else, even if they have the same condition as you.

Things to be careful of

When treatment with this product Should be used with caution in patient with impaired renal function. Please strictly adhere to the doctor's instructions.

If you are feeling abnormal or unusual problems please call your doctor.

Side effects

Common side effect are as follows:

- tiredness

- headache
- abdominal discomfort
- abdominal pain
- lung infection
- feeling sick (nausea)
- being sick (vomiting)
- diarrhea
- muscle pain and discomfort
- joint pain

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my

(Consumers→Reporting)

Storage and Disposal of VALFIX F.C. TABLET

Storage

Keep in a tight container. Store below 30°C, protect from light & moisture.


Disposal

Tell your doctor or pharmacist if you stop taking the tablets or the tablets have passed their expiry date. This medication should not be disposed of via wastewater or household waste. Ask them what to do with any tablets that are left over.

Product description

What it looks like

An orange-brown elliptical tablet, one side impressed with 'BM'.



Ingredients

Active(s):-
Lamivudine

Non active ingredients:-
Microcrystalline Cellulose
Sodium Starch Glycolate
Magnesium Stearate
Colloidal Silicon Dioxide
Hydroxyl Methylcellulose
Polyethylene Glycol
Titanium Dioxide
Iron Oxide Yellow
Iron Oxide Red

MAL no.

MAL

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13, Kawasan
Perindustrian Bandar Baru Bangi, 43000
Kajang, Selangor D.E., Malaysia.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan
Malaysia.

Date of revision

07/11/2016