

SHINALLERG EYE DROP

Tetrahydrozoline HCl, Antazoline HCl (0.4mg/ml, 0.5mg/ml)

What is in this leaflet

This leaflet contains information below:

1. What SHINALLERG EYE DROP is used for.
2. How SHINALLERG EYE DROP works.
3. Before you use SHINALLERG EYE DROP.
4. How to use SHINALLERG EYE DROP.
5. While you are using SHINALLERG EYE DROP.
6. Side effects.
7. Storage and Disposal of SHINALLERG EYE DROP.
8. Product description
9. Manufacturer & Product Registration Holder
10. Date of revision

What SHINALLERG EYE DROP is used for

It is used to relieve allergic reaction/symptoms (eg. redness, swelling, itching) in eye.

How SHINALLERG EYE DROP works

Antazoline is anti-histamine which provides relief of itching. Tetrahydrozoline is a sympathomimetic which reduces redness and swelling in the eyes. When both are used together, they are able to relieve symptoms of allergic eye disease effectively.

Before you use SHINALLERG EYE DROP

When you must not apply it

If you are allergic to antazoline and/or tetrahydrozoline or to any of the ingredients of SHINALLERG EYE DROP. If you wear contact lenses, they should be removed before instillation and not reinserted for at least 15 minutes.

Before you start to apply it

Tell your doctor if :

- You are elderly and have high blood pressure, a heart condition, a thyroid problem or diabetes.
- You are pregnant or breastfeeding.
- You suffer from dry eyes.
- You have an adrenal tumour (phaeochromocytoma).

Taking other medicines

Concomitant use of SHINALLERG EYE DROP with Monoamine Oxidase Inhibitors (MAOI) is prohibited. Tell your doctor if you use any other eye drops.

How to use SHINALLERG EYE DROP

How much to apply

Apply 1 drop every three hours during the acute phase, for maintenance therapy 1 drop 2-3 times per day is sufficient. 1-2 drops per day are sufficient for children.

When to apply it

SHINALLERG EYE DROP should be applied according to doctor's instruction.

How long to apply it

SHINALLERG EYE DROP should be applied according to doctor's instruction. Do not use for more than 14 days without consulting a doctor.

If you forgot to apply it

If you miss a dose of SHINALLERG EYE DROP and you are applying it regularly, apply it as soon as possible. If several hours have passed or if it is nearing time for the next dose, do not double the dose to catch up, unless advised by your health care provider. Do not apply two doses at once.

If you apply too much

Quickly seek for medical attention if you think that you or anyone else has applied too much SHINALLERG EYE DROP.

While you are using SHINALLERG EYE DROP

Things you must do

Apply the eye drops according to doctor's instruction.

Things you must not do

If you wear contact lenses, they should be removed before instillation and not reinserted for at least 15 minutes. Do not stop applying this medicine or change the dosing regimen without instruction from your doctor. Do not give SHINALLERG EYE DROP to anyone else, even if they have the same condition as you.

Things to be careful of

Use with caution in elderly patients with severe cardiovascular disease, including arrhythmia, poorly controlled hypertension or diabetes. If symptoms persist for 3 to 4 days, you should consult the doctor.

Side effects

SHINALLERG EYE DROP may cause drowsiness, dizziness, sleepiness, sweating, tremor, headache, central excitation or blurred vision. Patient may experience burning / stinging upon instillation, which is mild and transient in nature. Rare side effects include chest pain, high blood pressure, increase in heart rate and hypersensitivity.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of SHINALLERG EYE DROP

Storage

Store at temperature below 30°C.

Disposal

SHINALLERG EYE DROP should be disposed when your doctor or pharmacist tells you to stop using this medicine or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required.

Discard this eye drops one month after open.

Product description

What it looks like

A clear and colorless solution.

Ingredients

Active Ingredient

Tetrahydrozoline Hydrochloride
Antazoline Hydrochloride

Inactive Ingredient

Benzalkonium Chloride
Sodium chloride
Hydroxypropyl Methylcellulose
Hydrochloric acid
Water for injection

MAL number

MAL17107026AZ

**Manufacturer and Product Registration
Holder**

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Date of revision:

28/02/2025

Serial Number:

NPRA(R2/1)250225/1421