

URSAPHARM			
Kunde:	Export / Pharmaforte Malaysia	Lackierung:	/
Produkt:	PB ALLERGO-COMOD 10 ML ATRO MY	Lackaussparung:	☐ /
Art.-Nr.:	90.1608	Pharmacode:	1541
Format:	185 x 275 mm (Plano)	Zusatzspezifikation:	/
Blindenschrift (f5):	JA / NEIN	Sonstiges:	/
Material:	Offset / 60 g/qm (UPM fine, Fa. Nordland)	Kleinste Schriftgröße:	10 Pt
Kleber (ET):	/	Farbtoleranz vorhanden:	JA / NEIN Nr.: FTK_UPH_
Farbe 1:	☒ Schwarz (1/1-fbg.)	Farbtoleranz fertigen:	JA / NEIN Nr.: FTK_UPH_
Farbe 2:	☐ /	(6-fache Ausfertigung)	
Farbe 3:	☐ /		
Farbe 4:	☐ /	Korrekturabzug Nr.:	1 (mock-up)
Farbe 5:	☐ /	Erstellt:	05.10.2021 C. Staab
Farbe 6:	☐ /	Freigabe Kunde:	
Farbe 7:	☐ /	Endfreigabe/Gut zum Druck:	

ALLERGO-COMOD® EYE DROPS

Sodium Cromoglycate 20 mg/ml
Anti-allergic agent

PRODUCT DESCRIPTION

Allergo-COMOD® eye drops is a clear and colourless to slightly yellow solution available in dropper plastic bottle with 10 ml solution.

PHARMACOLOGY

Pharmacodynamics & Pharmacokinetics

Allergo-COMOD® eye drops contain an anti-allergic agent which acts by inhibiting the degranulation of mast cells. By this, the release histamine is suppressed.

INDICATIONS AND CLINICAL USE


Indicated in acute and chronic allergic conjunctivitis including hay-fever conjunctivitis, vernal keratoconjunctivitis.

DOSAGE AND ADMINISTRATION

Instill 1 drop 4 times daily into the conjunctival sac. Children 4 years and above and adults apply the same dosage. The exact duration of therapy should be based on the physician’s judgement.

Remove the cap covering the nozzle. Before applying Allergo-COMOD® eye drops for the first time, press on the bottom of the bottle as frequently and necessary until the first drops is coming out. Afterwards the bottle is ready for all following applications.

Invert the bottle as is shown in the picture and press quickly and powerful on the bottom of the bottle activating the mechanism to get out one drop.



Because of the special construction of valves the outcoming drops are always of equal size and speed not depending on how powerful you press the bottom.

Tilt your head back, with a finger retract the lower lid until the inner red, moist surface is exposed and bring the eye drops as described into the conjunctival sac. Close the eyes slowly. After use close the bottle carefully. A contact of the bottle with eye or skin should be avoided.

The treatment with Allergo-COMOD® eye drops should not be stopped after symptoms improve. Rather its use should be continued until the patient is no longer in contact with the allergen (pollen, house dust, mold, etc.).

MODE OF ADMINISTRATION

For ophthalmic use only.

CONTRAINDICATIONS

In hypersensitivity against one of the ingredients.
Contact lenses should not be worn during treatment with Allergo-COMOD® eye drops.

WARNINGS AND PRECAUTIONS

Immediately after applying Allergo-COMOD® eye drops vision will be blurred for a short time which subsequently impair reaction time while driving or operating machinery. Safety and effectiveness in children below 4 years of age has not been established.

DRUG INTERACTIONS

Not known.

PREGNANCY AND LACTATION

Although there is no indication to teratogenic effects, Allergo-COMOD® eye drops is not recommended for use during the first three month of pregnancy.

ADVERSE REACTIONS

In rare cases temporary burning, foreign body-sensation, hyperemia and chemosis have been reported.

OVERDOSAGE

Overdosing or intoxications are not to be expected when Allergo-COMOD® eye drops are used properly. Until now, no cases of intoxications are known.

STORAGE CONDITIONS

Do not use Allergo-COMOD® eye drops longer than 12 weeks after the bottle has been opened.
Do not store above 30°C.

AVAILABILITY OF DOSAGE FORMS

10 mL

AVAILABLE PACK SIZE

1 x 10 mL

MANUFACTURER

Ursapharm Arzneimittel GmbH
66129 Saarbrücken/Germany

DATE OF REVISION

October 2021

