

Gemita 1 g Label Malaysia - Unit-II

Review 1

140%

110 x 32 mm

Each vial contains:
 Gemcitabine Hydrochloride USP
 equivalent to Gemcitabine
 Mannitol USP

Store below 30°C. Do not refrigerate.

SINGLE DOSE VIAL

FOR I.V. INFUSION AFTER DILUTION.

Keep out of the reach of children.
 Jauhkan dari kanak-kanak.

Dosage: As directed by the Physician. Read accompanying package insert for detailed dosage, directions for use and precautions.

Direction: Reconstitute Gemcitabine for Injection USP 1 g with 25 ml of Sterile 0.9% w/w Sodium Chloride Injection. Shake gently to make a clear solution. This dilution yields a gemcitabine concentration of 38 mg/ml, which includes accounting for the displacement volume of the lyophilized powder (1.3 ml for 1 g vial). The reconstituted solution should be used immediately or within 24 hours if stored **below 30°C**.

Discard unused portion.

Warning: Cytotoxic agent.

1 g

R

Gemcitabine

for Injection USP

Gemita

LYOPHILIZED

Caution: It is dangerous to take this preparation except under Medical Supervision.

UBAT TERKAWAL CONTROLLED MEDICINE

Mfg. Lic. No.: MB/07/519

Registration No.: MAL08042477AZ

Product Registration Holder:
Fresenius Kabi Malaysia Sdn Bhd
 3-1 & 3-2, Axis Technology Centre, Lot 13,
 Jalan STA/225, 46100 Petaling Jaya, Selangor,
 Malaysia. Tel: 03-79579292

Manufactured by: **Fresenius Kabi Oncology Limited**,
 Village Krishnapura, Baddi, Tehsil Nalagarh, District Solan,
 IN-174101 India

BATCH:

MFG:

EXP:



	Pantone 285 C
	Pantone 7413 C
	Pantone 300 C
	Pantone 1795 C
	Black
	Unvarnish area (38 x 12 mm)