


Batch No.:
Manuf. Date :
Exp. Date:



SmofKabiven Peripheral

Emulsion for infusion

800 kcal

1206 ml

PERIPHERAL

 **FRESENIUS KABI**

Intravenous use, infusion into a peripheral or central vein. For single infusion only.

Energy content total approx. 800 kcal
Energy content non-protein approx. 700 kcal

Corresponding to:

Carbohydrates (glucose anhydrous):	85 g
Amino acids:	38 g
Nitrogen:	6.2 g
Lipids:	34 g

Electrolytes;

Sodium:	30 mmol
Potassium:	23 mmol
Magnesium:	3.8 mmol
Calcium:	1.9 mmol
Phosphate (contribution from lipid emulsion and amino acid solution):	9.9 mmol
Zinc:	0.03 mmol
Sulphate:	3.8 mmol
Chloride:	27 mmol
Acetate:	79 mmol

Osmolality: approx. 950 mosmol/kg water
Osmolarity: approx. 850 mosmol/l
pH: approx. 5.6

A three chamber bag (1206 ml) contains:

1. Glucose 13 %: Glucose (as monohydrate) 85 g.

2. Amino acid solution with electrolytes: Alanine 5.3 g, arginine 4.6 g, glycine 4.2 g, histidine 1.1 g, isoleucine 1.9 g, leucine 2.8 g, lysine (as acetate) 2.5 g, methionine 1.6 g, phenylalanine 1.9 g, proline 4.2 g, serine 2.5 g, taurine 0.38 g, threonine 1.7 g, tryptophan 0.76 g, tyrosine 0.15 g, valine 2.4 g, calcium chloride (as dihydrate) 0.21 g, sodium glycerophosphate (as hydrate) 1.6 g, magnesium sulphate (as heptahydrate) 0.46 g, potassium chloride 1.7 g, sodium acetate (as trihydrate) 1.3 g, zinc sulphate (as heptahydrate) 0.005 g.


3. Lipid emulsion: Refined soya-bean oil 10.2 g, medium-chain triglycerides 10.2 g, refined olive oil 8.5 g, fish oil rich in omega-3-acids 5.1 g.

Excipients: Glycerol, purified egg phospholipids, all-*rac*- α -tocopherol, sodium hydroxide (pH adjuster), sodium oleate, acetic acid glacial (pH adjuster), water for injections.

Read the package leaflet before use.
Keep out of the reach and sight of children.

The contents of the three separate chambers have to be mixed before use. Additions should be made aseptically.

After mixing the product should be used immediately.
For single infusion only.



Do not store above 25°C. Do not freeze. Store in overpouch.
Any unused solution remaining after infusion should be discarded.

Approved 2019-01-16:


Regulatory Affairs Manager
ADDITIVE PORT


INFUSION PORT

Approx.


1000

750

500

250






337 658 (ersätter 332 903) 4:e korr 19-01-16, M Eriksson, 018 64 40 46

Format: 320x 350 mm (2a)

Batch No.:
Manuf. Date :
Exp. Date:



SmofKabiven Peripheral

Emulsion for infusion

1000 kcal
1448 ml

Intravenous use, infusion into a peripheral or central vein. For single infusion only.

PERIPHERAL



Energy content total approx. 1000 kcal
Energy content non-protein approx. 800 kcal

Corresponding to:

Carbohydrates (glucose anhydrous):	103 g
Amino acids:	46 g
Nitrogen:	7.4 g
Lipids:	41 g


Electrolytes;

Sodium:	36 mmol
Potassium:	28 mmol
Magnesium:	4.6 mmol
Calcium:	2.3 mmol
Phosphate (contribution from lipid emulsion and amino acid solution):	11.9 mmol
Zinc:	0.03 mmol
Sulphate:	4.6 mmol
Chloride:	32 mmol
Acetate:	96 mmol

Osmolality: approx. 950 mosmol/kg water
Osmolarity: approx. 850 mosmol/l
pH: approx. 5.6

A three chamber bag (1448 ml) contains:

- Glucose 13 %:** Glucose (as monohydrate) 103 g.
 - Amino acid solution with electrolytes:** Alanine 6.4 g, arginine 5.5 g, glycine 5.1 g, histidine 1.3 g, isoleucine 2.3 g, leucine 3.3 g, lysine (as acetate) 3.0 g, methionine 1.9 g, phenylalanine 2.3 g, proline 5.1 g, serine 3.0 g, taurine 0.46 g, threonine 2.0 g, tryptophan 0.91 g, tyrosine 0.17 g, valine 2.9 g, calcium chloride (as dihydrate) 0.26 g, sodium glycerophosphate (as hydrate) 1.9 g, magnesium sulphate (as heptahydrate) 0.55 g, potassium chloride 2.0 g, sodium acetate (as trihydrate) 1.6 g, zinc sulphate (as heptahydrate) 0.006 g.
 - Lipid emulsion:** Refined soya-bean oil 12.3 g, medium-chain triglycerides 12.3 g, refined olive oil 10.1 g, fish oil rich in omega-3-acids 6.1 g.
- Excipients:** Glycerol, purified egg phospholipids, all-*rac*- α -tocopherol, sodium hydroxide (pH adjuster), sodium oleate, acetic acid glacial (pH adjuster), water for injections.



Read the package leaflet before use.
Keep out of the reach and sight of children.

The contents of the three separate chambers have to be mixed before use. Additions should be made aseptically.

After mixing the product should be used immediately.
For single infusion only.

Do not store above 25°C. Do not freeze. Store in overpouch.
Any unused solution remaining after infusion should be discarded.

Manufacturer:

Fresenius Kabi AB,
751 74 Uppsala, Sweden

MAL11080004ARZ

Jauhi daripada kanak kanak.

Controlled Medicine/Ubat Terkawal

Product Registration Holder/

Importer:

Fresenius Kabi Malaysia Sdn Bhd
3-1 & 3-2, Axis Technology Centre,
Lot 13, Jalan 51A/225,
46100 Petaling Jaya,
Selangor, Malaysia.

337 659

approx.

1000

750

500

250

Approved 2019-01-16:

Anne Strand
Regulatory affairs manager


ADDITIVE PORT



INFUSION PORT



Batch No.:
Manuf. Date :
Exp. Date:



SmofKabiven Peripheral

Emulsion for infusion

1300 kcal
1904 ml

Intravenous use, infusion into a peripheral or central vein. For single infusion only.

PERIPHERAL



**FRESENIUS
KABI**

Energy content total approx.
1300 kcal
Energy content non-protein
approx. 1100 kcal

Corresponding to:

Carbohydrates (glucose anhydrous):	135 g
Amino acids:	60 g
Nitrogen:	9.8 g
Lipids:	54 g


Electrolytes;

Sodium:	48 mmol
Potassium:	36 mmol
Magnesium:	6.0 mmol
Calcium:	3.0 mmol
Phosphate (contribution from lipid emulsion and amino acid solution):	15.6 mmol
Zinc:	0.05 mmol
Sulphate:	6.1 mmol
Chloride:	42 mmol
Acetate:	125 mmol

Osmolality: approx. 950 mosmol/
kg water
Osmolarity: approx. 850
mosmol/l
pH: approx. 5.6

A three chamber bag (1904 ml) contains:

- Glucose 13 %:** Glucose (as monohydrate) 135 g.
- Amino acid solution with electrolytes:** Alanine 8.4 g, arginine 7.2 g, glycine 6.6 g, histidine 1.8 g, isoleucine 3.0 g, leucine 4.4 g, lysine (as acetate) 4.0 g, methionine 2.6 g, phenyl-alanine 3.1 g, proline 6.7 g, serine 3.9 g, taurine 0.60 g, threonine 2.6 g, tryptophan 1.2 g, tyrosine 0.24 g, valine 3.7 g, calcium chloride (as dihydrate) 0.34 g, sodium glycerophosphate (as hydrate) 2.5 g, magnesium sulphate (as heptahydrate) 0.72 g, potassium chloride 2.7 g, sodium acetate (as trihydrate) 2.0 g, zinc sulphate (as heptahydrate) 0.008 g.
- Lipid emulsion:** Refined soya-bean oil 16.1 g, medium-chain triglycerides 16.1 g, refined olive oil 13.4 g, fish oil rich in omega-3-acids 8.0 g.



Excipients: Glycerol, purified egg phospholipids, all-*rac*- α -tocopherol, sodium hydroxide (pH adjuster), sodium oleate, acetic acid glacial (pH adjuster), water for injections.

Read the package leaflet before use.
Keep out of the reach and sight of children.

The contents of the three separate chambers have to be mixed before use. Additions should be made aseptically.

After mixing the product should be used immediately.
For single infusion only.

Do not store above 25°C. Do not freeze. Store in overpouch.

Any unused solution remaining after infusion should be discarded.

Manufacturer:

Fresenius Kabi AB,
751 74 Uppsala, Sweden

MAL11080004ARZ
Jauhi daripada kanak kanak.
Controlled Medicine/Ubat Terkawal
Product Registration Holder/Importer:
Fresenius Kabi Malaysia Sdn Bhd
3-1 & 3-2, Axis Technology Centre,
Lot 13, Jalan 51A/225,
46100 Petaling Jaya,
Selangor, Malaysia.

337 660

Approx.
1500

1000

750

500



ADDITIVE PORT



INFUSION PORT

Approved 2019-01-16:

Janne Strand

Regulatory affairs manager