

TERBRON SYRUP

Terbutaline Sulphate (1.5 mg/5ml)

What is in this leaflet

1. What *Terbron syrup* is used for
2. How *Terbron syrup* works
3. Before you use *Terbron syrup*
4. How to use *Terbron syrup*
5. While you are using it
6. Side effects
7. Storage and Disposal of *Terbron syrup*
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What *Terbron syrup* is used for

For the relief of bronchospasm (constriction of the airways going into the lungs) in bronchial asthma and bronchitis (inflammation in the main air passages).

How *Terbron syrup* works

The active ingredient in this medicine is Terbutaline Sulphate. Terbutaline is a bronchodilator (beta-2 receptor agonist) that produces relaxation of bronchial smooth muscle.

Before you use *Terbron syrup*

- When you must not use it

- If you are allergic (hypersensitive) to sympathomimetic amines or any of the other ingredients of *Terbron syrup*.
- If you are patients with thyrotoxicosis (excess thyroid hormone in the body).

- Before you start to use it

Check with your doctor or pharmacist before using your medicine if:

- You have heart diseases.
- You have diabetes mellitus.
- You have hyperthyroidism (overactive thyroid gland).
- You have high blood pressure.

Pregnancy and lactation

Do not use *Terbron syrup* if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not use *Terbron syrup* if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

- Taking other medicines

There are a few medicines which may interact with *Terbron syrup*. It is important to tell your doctor or pharmacist about all the medicines that you are taking, including those obtained without prescription. You should talk to your doctor or pharmacist if you are taking:

- Sympathomimetic agent (treat cardiac arrest and low blood pressure)
- Beta-blocker (reduce blood pressure)
- Monoamine oxidase inhibitors and tricyclic antidepressants

How to use *Terbron syrup*

- How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Adults:

1 tablespoonful, 2-3 times daily

Children:

Below 3 years: ½ teaspoonful, 3 times daily

3-7 years: ½ teaspoonfuls, 3 times daily

7-15 years: 1-2 teaspoonfuls, 3 times daily

Or as prescribed by the physician.

- When to use it

Use *Terbron syrup* when needed and according to doctor prescription.

- How long to use it

Continue taking *Terbron syrup* or as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking *Terbron syrup*.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give *Terbron syrup* to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Be careful when engaging in activities requiring mental alertness such as driving a car or operating machinery until you know how this medicine affects you.

Side effects

Like all medicines, *Terbron syrup* can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

The side effects are an increase in heart rate, changes in blood pressure, nervousness, tremor, palpitation and dizziness.

Headache, nausea, vomiting, anxiety, restlessness, lethargy, drowsiness, weakness, flushes, sweating, chest

discomfort, muscle cramps and tinnitus (perception of noise or ringing in the ears) have also been reported.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Date of revision

05/01/2021

Serial Number

NPRA(R2/1)190312/82

Storage and Disposal of Terbron syrup

- Storage

Keep out of the reach and sight of children.

Store at temperature not exceeding 30°C, protect from light and excessive heat.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Clear, almost colorless, cherry-flavored solution.

- Ingredients

- Active ingredient
Terbutaline sulphate

- Inactive ingredients
Sodium citrate dihydrate, citric acid monohydrate, sucrose, propylene glycol, methylparaben, propylparaben, sorbitol solution 70%, cherry cider essence, purified water

- MAL number):

MAL19900433AZ

Manufacturer

Biolab Company Ltd
625 Soi, 7A Bangpoo Industrial Estate,
Samutprakarn 10280,
Thailand

Product Registration Holder

Medispec (M) Sdn Bhd
No. 55 & 57, Lorong Sempadan 2
11400 Ayer Itam, Penang, Malaysia