

MOTIGUT SUSPENSION

Domperidone (1 mg/ml)

What is in this leaflet

1. What *Motigut Suspension* is used for
2. How *Motigut Suspension* works
3. Before you use *Motigut Suspension*
4. How to use *Motigut Suspension*
5. While you are using it
6. Side effects
7. Storage and Disposal of *Motigut Suspension*
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What *Motigut Suspension* is used for

Motigut is used for the relief of symptoms of nausea and vomiting. This includes:

- Nausea and vomiting of functional, organic, infectious or dietary origin.
- Nausea and vomiting induced by:
 - Radiotherapy or drug therapy
 - Dopamine agonists (such as L-dopa and bromocriptine) used in the treatment of Parkinson's disease.

How *Motigut Suspension* works

Motigut suspension contains domperidone with anti-emetic (stop vomiting) properties. It also increases the gastric emptying.

Before you use *Motigut Suspension*

- When you must not use it

Do not use *Motigut suspension* if:

- You are allergic to domperidone or any of the ingredients in this medication.
- You suffer from a tumor of the pituitary gland called prolactinoma
- You have or had heart problems including heart failure or heart rhythm disorders including a slow or irregular heartbeat
- You are taking QT-prolonging drugs
- You are taking potent CYP3A4 inhibitors
- • have bleeding, obstruction and hole in the stomach and bowel (e.g. severe stomach cramps or persistent black stools).
- You have moderate to severe liver disease

Do not take *Motigut Suspension* if you are taking any of the following medicine: Clarithromycin (antibiotic)

Pregnancy and lactation

Do not take *Motigut Suspension* if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take *Motigut Suspension* if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start to use it

Tell your doctor or pharmacist if:

- It is for infant or small children
- You have liver problems
- You have kidney problems
- You have heart disease
- You have low or high potassium level or low magnesium level.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

It is especially important to tell your doctor or pharmacist if you are taking any of the following:

- CYP3A4 inhibitors (eg: ketoconazole (antifungal), ritonavir (antiviral) and erythromycin (antibiotic))
- Medicines that affect the ECG results

How to use *Motigut Suspension*

- How much to use

You should always take the lowest amount of *Motigut Suspension* that works for you and you should not take it for longer than is necessary. Although the amount of *Motigut Suspension* you should usually take is described below, your doctor may adjust your dose to your personal needs.

Motigut Suspension is most effective if taken 15-30 minutes before meals.

Adults and adolescents (12 years of age and over) weighing 35 kg or more; children weighing 35kg or more:

Tablets: Take 1 tablet 3 to 4 times a day. Do not take more than 4 tablets per day (40 mg/day).

Oral suspension: Take 10mL of oral suspension 3 or 4 times a day. Do not take more than 40mL per day (40 mg/day).

Adults and adolescents weighing less than 35 kg:

Oral suspension: Give 0.25 milliliters of the oral suspension per kilogram of body weight 3 or 4 times a day. The maximum dose per day is 1 mg/kg but do not exceed 35 mg per day.

Infants and children less than 12 years of age and weighing less than 35kg:

The effectiveness of *Motigut Suspension* has not been established in infants and children under 12 years of age with a body weight of <35 kg.

- When to use it

To be taken 3-4 times a day or as directed by your doctor or pharmacist.

For people with kidney problems, the dosing frequency of *Motigut* should be reduced to once or twice daily.

It is recommended to take *Motigut* 15-30 minutes before meals.

- How long to use it

Continue taking *Motigut Suspension* for as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause drowsiness, disorientation and extrapyramidal reaction (involuntary muscle movements, loss of movement especially in children).	extrapyramidal effects, transient intestinal cramps, hives, breast enlargement, absence of menstrual period and others.	- <u>MAL number:</u> MAL10120032AZ
While you are using it - <u>Things you must do</u> Take your medicine exactly as your doctor has told you.	Very rare side effects include abnormal heartbeat, ECG changes, and sudden loss of heart function	Manufacturer Square Pharmaceuticals PLC. Square Road, Salgaria, Pabna,6600, Bangladesh.
Tell all the doctors, dentists and pharmacists treating you that you are taking <i>Motigut Suspension</i>.	You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumer→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].	Product Registration Holder Medispec (M) Sdn Bhd B-1-07, Jalan SS 25/22, Taman Mayang, 47301 Petaling Jaya, Selangor, Malaysia.
Tell your doctor immediately if you become pregnant while taking this medication.	Storage and Disposal of <i>Motigut Suspension</i>	Date of revision 09/09/2024
- <u>Things you must not do</u> Do not stop taking the medicine unless advised by your doctor.	- <u>Storage</u> Keep out of the reach and sight of children.	Serial number NPRA(R4/1)240118/00022
Do not take any new medicines without consulting your doctor or pharmacist.	Store below 30 °C.	
Do not give <i>Motigut Suspension</i> to anyone else, even if they have the same symptoms or condition as you.	- <u>Disposal</u> Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.	
- <u>Things to be careful of</u> <i>Driving and using machines</i> This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.	Product Description	
The oral suspension contains sorbitol and may be unsuitable for people with sorbitol intolerance.	- <u>What it looks like</u> A white raspberry-banana flavored suspension having sweet taste, filled and sealed in amber PET bottle with white color SQUARE printed cap.	
Stop taking domperidone and consult your physician if you have irregular heart beat	- <u>Ingredients</u> - Active ingredient Domperidone	
Side effects Like all medicines, <i>Motigut Suspension</i> can cause side effects, although not everybody gets them.	- Inactive ingredients Liquid Sorbitol (non-crystalizing) Sodium methyl hydroxybenzoate Dispersible cellulose Sodium propyl hydroxybenzoate Saccharin sodium Carmellose sodium Polysorbate 60 Anhydrous citric acid Sodium hydroxide Banana liquid essence Raspberry liquid essence Purified water	
Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.		
Side effects may include allergic reaction, increase milk production,		