

BELCID SUSPENSION

Aluminum Hydroxide/Magnesium Hydroxide/Simethicone (918mg/300mg/60mg per 15ml)

What is in this leaflet

1. What BELCID SUSPENSION is used for
2. How BELCID SUSPENSION works
3. Before you use BELCID SUSPENSION
4. How to use BELCID SUSPENSION
5. While you are using it
6. Side effects
7. Storage and Disposal of BELCID SUSPENSION
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What BELCID SUSPENSION is used for

BELCID SUSPENSION is used in symptomatic relief of gastric hyperacidity, heartburn, flatulence (wind) and stomach and intestinal ulcer pain.

How BELCID SUSPENSION works

BELCID SUSPENSION contains Aluminum hydroxide, Magnesium hydroxide and Simethicone. It is an antacid which relieves symptoms by neutralising stomach acid and has an antifoaming agent which aids removal of gas.

Before you use BELCID SUSPENSION

- When you must not use it

Do not take BELCID SUSPENSION if:

- You are allergic to any of the active and other ingredients.
- You suffer from kidney failure.

- Before you start to use it

Ask your pharmacist or doctor before taking BELCID SUSPENSION if:

- You have kidney insufficiency.

- Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Especially:

- Tetracycline (antibiotic)
- Digoxin (for heart disease)
- Indomethacin (pain killer)
- Iron salts (iron supplement)

- Dicumarol (blood thinning).

How to use BELCID SUSPENSION

- How much to use

1 tablespoonful, 4 times daily, 1 hour after meals and at bedtime or as directed by the physician.

- When to use it

BELCID SUSPENSION should not be taken at the same time as other medicines. These medicines should be spaced 1-2 hours from doses of BELCID.

- How long to use it

Take BELCID SUSPENSION exactly as your doctor has told you. If you are not sure, check with your doctor or pharmacist.

- If you forget to use it

If you forget to take a dose, skip the missed dose. Then take your next dose as normal.

- If you use too much (overdose)

If you take more BELCID SUSPENSION than you should, tell your doctor or go to hospital casualty department straight away.

While you are using it

- Things you must do

- If symptoms persist consult your doctor.
- Shake well before use.

- Things you must not do

- Do not allow this product to freeze.
- Do not take BELCID SUSPENSION after the expiry date printed on the pack.

- Things to be careful of

If you are pregnant, planning to become pregnant or breast feeding tell your doctor or pharmacist before taking this medicine.

Side effects

Tell your doctor or pharmacist if you notice any other unwanted effects.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-

78835550, or visiting the website nppra.moh.gov.my (Public→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI))

Storage and Disposal of BELCID SUSPENSION

- Storage

Store in dry place at temperature not exceeding 30°C.

Keep medicine out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

White with peppermint flavored suspension.

- Ingredients

- Active ingredients

Aluminum Hydroxide, Magnesium Hydroxide and Simethicone

- Inactive ingredients

Sorbitol Solution 70%, Saccharin Sodium, Peppermint Oil, Methylparaben, Propylparaben, Guar gum and Purified Water

- MAL number:

MAL19911699X

Manufacturer

Biolab Co. Ltd
625 Soi, 7A ,Bangpoo Industrial Estate,
Samutprakarn 10280, Thailand

Product Registration Holder

Medispec (M) Sdn.Bhd
No. 55 & 57, Lorong Sempadan 2,
(Off Boundary Road),
11400 Ayer Itam, Penang,
Malaysia.

Date of revision

10.8.2018

Serial Number:

NPRA(R4/1)180118/00019 at the end of the RiMUP