

MUCOSTOP 600 mg Effervescent Tablet

Acetylcysteine (600 mg)

What is in this leaflet

1. What **MUCOSTOP** is used for
2. How **MUCOSTOP** works
3. Before you use **MUCOSTOP**
4. How to use **MUCOSTOP**
5. While you are using it
6. Side effects
7. Storage and Disposal of **MUCOSTOP**
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What MUCOSTOP is used for

MUCOSTOP is used for treatment of respiratory diseases which leads to the formation of a lot of thick secretions which are difficult to be coughed out such as acute and chronic bronchitis (inflammation of the airways that carry air to your lungs), inflammation of the throat, sinus and trachea, flu infection, asthma and mucoviscidosis (abnormal mucus production in lungs).

How MUCOSTOP works

Acetylcysteine has strong fluidification activity to mucus and pus secretion. It acts by reducing inflammation and increasing mucosa regeneration speed.

Before you use MUCOSTOP

- When you must not use it

DO NOT USE MUCOSTOP in the following situations:

- If you are allergic to acetylcysteine or any of the other ingredients of MUCOSTOP
- In children below 2 years old
- If you are suffering from phenylketonuria
- If you have an active peptic ulcer.

Pregnancy and lactation

- Pregnancy category B.
- Not known whether acetylcysteine is distributed into milk.
- Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start to use it

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food

shop. Please also inform your doctor if you have asthma.

- Taking MUCOSTOP with other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use MUCOSTOP

- How much to use

Adults:

1 effervescent tablet a day (preferably in the evening)

- When to use it

Dissolve in water before administration.

- How long to use it

- Do not exceed the recommended dose.
- Your doctor may recommend a higher or lower dose, depending on the response to the treatment.
- Do not change the dose or stop treatment before talking to your doctor.

Administration way and method:

- MUCOSTOP is only used orally.
- After meal, dissolve the effervescent tablet in a glass of water and drink it straight away. Effervescent tablets should not be chewed and should not be swallowed.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are

no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause nausea, vomiting and diarrhea.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking MUCOSTOP.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give MUCOSTOP to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

- If you have gastric, it should be taken after food.
- If you have asthma, please inform your doctor before continuing this product.
- It is better not to mix other drugs with the MUCOSTOP solution.

Driving and using machines

MUCOSTOP has not been reported to have any adverse effects on vehicle and machine use.

Side Effects

As with all medicines, there may be side effects in people who are sensitive to the ingredients contained in MUCOSTOP.

MUCOSTOP may cause severe allergy and serious skin reactions. Stop using MUCOSTOP and seek medical assistance immediately if you experience any of the following symptoms:

- Severe allergy: breathing difficulties, light headedness, skin swellings or rash.

- Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

Other side effects reported with acetylcysteine include nausea, vomiting, fever, temporary loss of consciousness, sweating, pain in joint, blurred vision, disturbances of liver function.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npdra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of MUCOSTOP

-Storage

Keep MUCOSTOP in the places where children cannot see, cannot reach, and in their packaging.

Store at room temperature below 30°C and dry place.

Do not forget to close the cap of the tube after every use. Do not leave the cap of the tube open.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

White, round, flat-faced tablets.

Appearance of solution should be clear after dissolving in water.

- Ingredients

- Active ingredient(s)
Acetylcysteine

- Inactive ingredient(s)

Micronized monosodium citrate, sodium hydrogen carbonate, aspartame, polyvinylpyrrolidone, sodium chloride and lemon flavor

- MAL number(s):

MAL*****

Manufacturer

Neutec İlaç San. Tic. A.Ş.

1. OSB 1. Yol No: 3

Adpazarı / SAKARYA / TURKEY

Product Registration Holder

MEDISPEC (M) SDN BHD

55 & 57, Lorong Sempadan 2,

(Off Boundary Road)

11400 Ayer Itam, Penang,

Malaysia

Date of revision

14/10/2019

Serial number

NPRA (R2) 19/156