



Composition: Each vial contains: Tigecycline USP 50 mg Excipients q.s. Store below 30°C. Keep out of reach of children. Storage condition after reconstitution: Once reconstituted 5.0 mL of 0.2% Sodium Chloride Injection or 5% Dextrose Injection or Lactated Ringer's Solution for Injection, TYLIN may be stored at 25°C for up to 24 hours (up to 8 hours in the vial and remaining time in the IV bag). Alternatively, TYLIN may be refrigerated at 2°C to 8°C for up to 48 hours following immediate transfer of the reconstituted solution into the IV bag. Reconstituted solution must be transferred and further diluted for IV infusion. Jauhkan dari anak-anak. UBAT TERKAWAL.	TYLIN Tigecycline Powder for Solution for Infusion 50 mg / vial Lyophilized For I.V. infusion only Pack Size: 1 Vial Product Registration Holder Bioscience (S) Pte Ltd 8, Fair View, Unit 08-20/22, Simei Majang, 47021 Pandan Jaya, Singapore, Malaysia	Mfg. Lic. No. : G281499 Batch No. Mfg. Date Expiry Date NVZ Size - 10mm x 10mm Registration No: ML220801542 Manufactured by: PMILAD006199 Gulfic Biosciences Limited Unit-2, Survey No. 171/3, IV, No.3, Near GHI, Khalipora 396424, Naranat, Gujarat, India
--	---	---

Vial Label Size: 63 mm x 23 mm