

Paclitaxel Seacross 6 mg/ml

Concentrate for solution
for infusion

MAL*****

Manufactured by:

Sichuan Huiyu Pharmaceutical
Company Limited
Building 3, No. 333 Hanyang Road,
Shizhong District,
Neijiang, CN-641000, CHINA

Product Registration Holder:

B-1-07, Jalan SS 25/22,
Taman Mayang,
47301 Petaling Jaya,
Selangor, Malaysia

CAUTION: CYTOTOXIC AGENT

Paclitaxel Seacross 6 mg/ml

Concentrate for solution
for infusion

100 mg 16.7 ml

One 16.7 ml vial
Intravenous use
For single use only

Dilution required before use

CAUTION: CYTOTOXIC AGENT

Lot :

EXP:

MFD:



- Vial before opening: Store below 30°C.
- After opening before dilution: Chemical and physical in-use stability has been demonstrated for 28 days at 25°C following multiple needle entries and product withdrawals.
- From a microbiological point of view, once opened the product may be stored for a maximum of 28 days at 25°C. Other in-use storage times and conditions are the responsibility of the user.
- After dilution: After dilution the solution is for single use only. Chemical and physical in-use stability of the solution prepared for infusion has been demonstrated at 2°C to 8°C and at 25°C for 7 days when diluted in 5% dextrose solution, 5% dextrose and 0.9% sodium chloride solution and for 14 days when diluted in 0.9% sodium chloride injection.

**Keep out of the sight and
reach of children / Jauhkan
daripada capaian kanak-kanak**



CAUTION: CYTOTOXIC AGENT

Internal code

Paclitaxel Seacross 6 mg/ml

Concentrate for solution
for infusion



- From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless dilution has taken place in controlled and validated aseptic conditions
- 1 ml of concentrate for solution for infusion contains 6 mg paclitaxel. One 16.7 ml vial contains 100 mg of paclitaxel.
- Ethanol 392.63 mg/ml.

**Controlled Medicine/
Ubat Terkawal**

Read the package leaflet
before use

**For intravenous infusion after
appropriate dilution**

CAUTION: CYTOTOXIC AGENT