

URON Effervescent Granules

Sodium Bicarbonate/Tartaric Acid/Citric Acid/Sodium Citrate (1.76g/0.89g/0.72g/0.63g)

What is in this leaflet

1. What URON is used for
2. How URON works
3. Before you use URON
4. How to use URON
5. While you are using it
6. Side effects
7. Storage and Disposal of URON
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

What URON is used for

URON is used:

- For prevention of urates crystallisation in gout therapy.
- For relieving discomfort in mild urinary tract infections.
- For symptomatic relief of dysuria (painful or difficult urination).
- To enhance the action of certain antibiotics, especially sulphonamides.

How URON works

URON works as a urinary alkaliser by raising the pH of urine, making your urine less acidic. Maintaining alkaline urine may help dissolve uric acid stones.

Before you use URON

- When you must not use it

Do not take URON if you:

- have kidney failure or high sodium level in your blood
- are taking other medicines to treat urinary tract infections, such as hexamine mandelate or hexamine Hippurate
- have heart failure
- are taking quinolone antibiotics

- Before you start to use it

Check with your doctor before you start to take this medicine if:

- you are on a low sodium diet.

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-

feeding before using any medicine, including URON

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy from a pharmacy, supermarket or health food shop without a prescription.

The following medicines may interact with URON:

- tetracycline, ciprofloxacin, norfloxacin, ofloxacin, hexamine hippurate, hexamine mandelate (antibiotics)
- laxatives
- chlorpropamide (a diabetes medicine)
- lithium (medicine to treat bipolar disorder)
- salicylates (pain killers)
- amphetamines (medicine to treat attention deficit hyperactivity disorder - ADHD)
- antacids (medicine to treat heart burn).

How to use URON

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- How much to use

Dissolve 1 to 2 sachets in cold water.

- When to use it

Consume four times daily or as directed by your doctor or pharmacist.

- How long to use it

Use this medicine for 5 days or for the duration that your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do

not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the emergency department of your nearest hospital, if you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much URON may cause your body to become too alkaline (metabolic alkalosis).

While you are using URON

- Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are using URON.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not use any new medicines without consulting your doctor.

Do not give URON to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

If you are required to be on a low sodium diet, be aware that URON contains sodium (650mg per sachet).

Side Effects

Like all medicines, URON can cause side effects, although not everybody gets them.

Possible side effects with URON are as follow:

- mild laxative effect (causing loose stools or bowel movement)
- build-up of alkali in the body (systemic alkalosis) if used excessively
- high level of sodium in the blood

URON Effervescent Granules

Sodium Bicarbonate/Tartaric Acid/Citric Acid/Sodium Citrate (1.76g/0.89g/0.72g/0.63g)

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of URON

- Storage

Store below 30°C.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Appearance of powder: White to off white coloured, lemon flavoured granules.

Appearance of solution: Reconstituted solution is clear and lemon flavoured.

- Ingredients

- Active ingredient(s)
 - Sodium Bicarbonate BP
 - Tartaric Acid BP
 - Citric Acid Anhydrous BP
 - Sodium Citrate Anhydrous USP
- Inactive ingredients
 - Lemon flavour

- MAL number:

MAL*****

Manufacturer

Ind-Swift Limited
Off. NH-21, Village Jawaharpur, Tehsil Derabassi, District SAS Nagar (Mohali), Punjab-140507, India.

Product Registration Holder

Medispec (M) Sdn Bhd,
B-1-07, Jalan SS 25/22,
Taman Mayang,
47301 Petaling Jaya, Selangor,
Malaysia

Date of revision

28.04.2026

Serial Number

NPRA (R2) 26/064

URON Effervescent Granules

Sodium Bicarbonate/Tartaric Acid/Citric Acid/Sodium Citrate (1.76g/0.89g/0.72g/0.63g)
