

TAVIN-EM FILM COATED TABLETS

Tenofovir Disoproxil Fumarate and Emtricitabine (300mg/200mg)

What is in this leaflet

1. What TAVIN-EM Tablet is used for
2. How TAVIN-EM Tablet works
3. Before you use TAVIN-EM Tablet
4. How to use TAVIN-EM Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of TAVIN-EM Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What TAVIN-EM Tablet is used for

TAVIN-EM Tablet is indicated in combination with other antiretroviral agents for the treatment of HIV-1 in adults.

TAVIN-EM Tablet is indicated in combination with safer sex practices for pre-exposure prophylaxis to reduce the risk of sexually acquired HIV-1 infection in adults.

How TAVIN-EM Tablet works

TAVIN-EM Tablet contains two active substances, emtricitabine and tenofovir disoproxil. Both of these active substances are antiretroviral medicines which are used to treat HIV infection. Emtricitabine is a nucleoside reverse transcriptase inhibitor and tenofovir is a nucleotide reverse transcriptase inhibitor. However, both are generally known as NRTIs and they work by interfering with the normal working of an enzyme (reverse transcriptase) that is essential for the virus to reproduce itself.

Before you use TAVIN-EM Tablet

- When you must not use it

Pregnancy and lactation

Do not take TAVIN-EM Tablet if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take TAVIN-EM Tablet if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start to use it

Tell your doctor if you

- have liver problems including hepatitis B virus infection
- have kidney problems or receive kidney dialysis treatment
- have bone problems

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

- other medicines that contain tenofovir or emtricitabine
- medicines that contain lamivudine
- adefovir

Especially tell your healthcare provider if you take

- Didanosine
- Atazanavir
- Lopinavir with ritonavir

How to use TAVIN-EM Tablet

- If you are taking TAVIN-EM Tablet to reduce the risk of getting HIV, take TAVIN-EM Tablet every day, not just when you think you have been at risk of HIV infection.

- If you are taking TAVIN-EM Tablet to reduce the risk of getting HIV, do not stop taking TAVIN-EM Tablet or miss any doses. Stopping use of TAVIN-EM Tablet, or missing doses, may increase your risk of getting HIV infection. Do not stop taking TAVIN-EM Tablet without contacting your doctor.

- How much to use

The recommended dose is:

Adults: one tablet each day with or without food.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the

label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking TAVIN-EM Tablet for as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking TAVIN-EM Tablet.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

TAVIN-EM FILM COATED TABLETS

Tenofovir Disoproxil Fumarate and Emtricitabine (300mg/200mg)

Do not give **TAVIN-EM Tablet** to anyone else, even if they have the same symptoms or condition as you.

- Do not share or re-use needles or other injection equipment.
- Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.
- Do not have any kind of sex without protection. Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.

- Things to be careful of

Driving and using machines

The effects of the medicine on ability to drive and use machine is unknown.

Side effects

TAVIN-EM Tablet has the following effects:

The most common side effects of emtricitabine and tenofovir disoproxil fumarate tablets include:

- Diarrhoea
- Nausea

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of TAVIN-EM Tablet

- Storage

Keep out of the reach and sight of children.
Store below 30°C in a tightly closed container.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

White to off white, modified capsule shaped, film-coated tablets, debossed with “EM” on one side “144” on the other side of the tablet.

- Ingredients

- Active ingredient(s)
Emtricitabine
Tenofovir Disoproxil Fumarate
- Inactive ingredients
Microcrystalline Cellulose,
Croscarmellose sodium,
Pregelatinized starch, Magnesium Stearate, Isopropyl alcohol, Opadry AMB white 80W68912, Purified water

- MAL number(s):

MAL*****

Manufacturer

EMCURE PHARMACEUTICALS LIMITED
Plot No. P-2, I.T.B.T. Park, Phase II,
M.I.D.C. Hinjawadi,
Pune Maharashtra 411057
India

Product Registration Holder

Medispec (M) Sdn Bhd
B-1-07, Jalan SS 25/22,
Taman Mayang,
47301 Petaling Jaya, Selangor, Malaysia

Date of revision

13/05/2026

Serial number

NPRA (R2) 26/010