

REXOM™ HYDROCORTISONE NEOMYCIN CREAM

Hydrocortisone Acetate BP/ Neomycin Sulphate BP (0.5% w/w / 0.5% w/w)

What is in this leaflet

1. What REXOM™ Hydrocortisone Neomycin is used for
2. How REXOM™ Hydrocortisone Neomycin works
3. Before you use REXOM™ Neomycin Hydrocortisone
4. How to use REXOM™ Neomycin Hydrocortisone
5. While you are using it
6. Side effects
7. Storage and Disposal
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What REXOM™ Hydrocortisone Neomycin is used for

It is indicated for the treatment of allergic dermatitis and other inflammatory skin diseases when complicated by infection caused by organisms susceptible to neomycin including contact dermatitis, neurodermatitis, atopic dermatitis (allergic eczema), non-specific anogenital pruritus and seborrheic dermatitis.

How REXOM™ Hydrocortisone Neomycin works

REXOM™ Hydrocortisone Neomycin Cream contains 0.5% w/w Hydrocortisone Acetate BP and 0.5% w/w Neomycin which is absorbed systemically.

Hydrocortisone: Occlusion or use over extensive areas increases absorption of topical dosage forms. It is mostly metabolized in skin.

Neomycin: Neomycin is readily absorbed from large denuded, burned or granulating areas and may cause nephrotoxicity and moderate to severe ototoxicity especially if given concurrently. The use of topical antibacterial may lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical or systemic use of the medication.

Before you use REXOM™ Hydrocortisone Neomycin**-When you must not use it**

- If you have tuberculous infection of skin, fungus infection, herpes simplex, vaccinia, varicella and extensive open wound.

-Before you start to use it

You must tell your doctor if:

- You have allergies to any medicine which you have taken previously to treat your current condition. .
- You are pregnant, trying to become pregnant or breast-feeding.

-Taking other medicines

Always inform your doctor and pharmacist if you are taking any other medicines, including herbal tonics, supplements and medicines that you buy without prescription.

How to use REXOM™ Hydrocortisone Neomycin

Always take this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. Do not take this medicine by mouth. This medicine is only for topical application.

-How much to use

Apply a thin layer to the affected skin area one to three times daily. When dilution is required, the diluted cream should not be used two weeks after issue.

-When to use it

Use as directed by your doctor or pharmacist.

-How long to take it

Continue using REXOM™ Hydrocortisone Neomycin for as long as your doctor recommends.

-If you forget to take it

- If you forget a dose, use it as soon as you remember it.
- However, if it is nearly time for the next dose, skip the missed one.
- Do not use a double dose to make up for a forgotten dose.

-If you use too much (overdose)

Symptoms of toxicity that is associated with topical corticosteroids therapy may be characterized as either local or systemic. Symptoms are described under 'Side effects'.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it**-Things you must do**

Tell your doctor if your symptoms worsen or they do not improve.

If you do become pregnant whilst using REXOM™ Hydrocortisone Neomycin tell your doctor.

-Things you must not do

Do not give REXOM™ Hydrocortisone Neomycin Cream to anyone else, even if their symptoms seem similar or they have same condition as you. Your doctor has prescribed REXOM™ Hydrocortisone Neomycin for you and your condition.

-Things to be careful of

Prolong use or use of large amount should be avoided in the treatment of skin treatment of skin infections following extensive burns tropic ulceration and other conditions where absorption of neomycin is possible.

Side effects

Like all medicines, REXOM™ Hydrocortisone Neomycin may cause side effects, although not everybody gets them. The following side effects may happen with this medicine.

- impaired wound healing
- thin fragile skin
- petechiae and ecchymoses
- facial erythema
- increased sweating
- suppress reactions to the skin tests may occur due to hydrocortisone
- Hypersensitivity reactions due to neomycin include itching, rashes, swelling or other sign of irritation not present before therapy.

REXOM™ HYDROCORTISONE NEOMYCIN CREAM

Hydrocortisone Acetate BP/ Neomycin Sulphate BP (0.5% w/w / 0.5% w/w)

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Consult your doctor or pharmacist if you are in doubt or for any further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website <http://npra.moh.gov.my> (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal

-Storage

Do not use after the expiry date stated on the label. The expiry date refers to the last day of that month.

Keep container tightly closed. Store below 30°C. Protect from freezing, heat and direct light.

Keep out of reach and sight of children.

-Disposal

Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

White to off-white colour, smooth, homogenous cream.

-Ingredients

• Active ingredient

Hydrocortisone Acetate BP (0.5% w/w),
Neomycin Sulphate BP (0.5% w/w)

• Inactive ingredients

Emulsifying Wax, Liquid Paraffin BP,
White Soft Paraffin BP, Methyl
Paraben, Propyl Paraben and Purified
Water.

MAL number

MAL19860242AZ

Manufacturer and Product

Registration Holder

KCK Pharmaceutical Industries Sdn.
Bhd.
Plot 61, Bayan Lepas Industrial Park
Phase 4, Lintang Bayan Lepas 1, 11900
Bayan Lepas, Pulau Pinang,
Malaysia.

Date of revision

26/10/2018

Serial Number

NPRA(R2/1)221107/972