

REXOM™ HYDROCORTISONE NEOMYCIN CREAM

Name and Strength of Active Ingredient(s) :

Hydrocortisone Acetate BP	0.5% w/w
Neomycin Sulphate BP	0.5% w/w
Preservatives :	
Methyl Paraben	0.15% w/w
Propyl Paraben	0.08% w/w

Product Description :

White to off-white colour, smooth, homogenous cream.

Pharmacodynamics :

Hydrocortisone : Occlusion or use over extensive areas increases absorption of topical dosage forms. It is mostly metabolized in skin.

Neomycin : Neomycin is readily absorbed from large denuded, burned or granulating areas and may cause nephrotoxicity and moderate to severe ototoxicity especially if given concurrently. The use of topical antibacterial may lead to skin sensitisation, resulting in hypersensitivity reactions with subsequent topical or systemic use of the medication.

Pharmacokinetics :

Hydrocortisone : It is absorbed systemically.

Neomycin : Neomycin is not absorbed through the intact skin. However, it is readily absorbed from large denuded, burned, or granulating areas.

Indication :

For the treatment of allergic dermatitis and other inflammatory skin diseases when complicated by infection caused by organisms susceptible to neomycin including contact dermatitis, neurodermatitis, atopic dermatitis (allergic eczema), non-specific anogenital pruritus and seborrhoeic dermatitis.

Recommended Dosage :

Apply a thin layer to the affected skin area one to three times daily. When dilution is required, the dilution cream should not be used two weeks after issue.

Mode of Administration :

Topical.

Contraindications :

In tuberculous infection of the skin, fungus infections, herpes simplex, vaccinia, varicella and in other cutaneous infections for which an infective antibiotic is not available for simultaneous application.

Warnings and Precautions :

Prolong use or use of large amount should be avoided in the treatment of skin infections following extensive burns, tropic ulceration and other conditions where absorption of neomycin is possible.

Interactions with Other Medicaments :

Not know Drug Interaction.

Pregnancy and Lactation :

Pregnancy Risk Factor C.

Adverse Effects :

Impaired wound healing, thin fragile skin, petechiae and ecchymosis, facial erythema, increased sweating and suppress reactions to skin tests may occur due to hydrocortisone. Hypersensitivity reactions due to neomycin include itching, rashes, swelling or other sign of irritation not present before therapy.

Symptoms and Treatment of Overdose :

Not available.

Storage Conditions :

Keep container tightly closed. Store below 30°C. Protect from freezing, heat and direct light.

Packaging Available :

15 gm tube, 15 gm x 25 tubes and 15gm x 50 tubes.
20 gm tub and 20gm x 10 tubs.

Registration Number :

MAL19860242AZ

Manufactured by & Product Registration Holder :

KCK PHARMACEUTICAL INDUSTRIES SDN. BHD.
Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision :

1 November 2023