

ANTAMAC SUSPENSION

Magnesium Trisilicate/Aluminium Hydroxide/Simethicone (500mg/250mg/25mg per 5ml)

What is in this leaflet

1. What Antamac is used for
2. How Antamac works
3. Before you use Antamac
4. How to use Antamac
5. While you are using Antamac
6. Side effects
7. Storage and disposal of Antamac
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Antamac is used for

Antamac is used in the symptomatic relief of acid indigestion, dyspepsia, heartburn and painful gastric symptoms.

How Antamac works

Aluminium and magnesium hydroxide, an antacid, neutralize or buffer existing quantities of stomach acid, thus providing relief of symptoms of excess acid. Simethicone, an antifatulent, relieves by belching or passing flatus.

Before you use Antamac

-When you must not use it

- Allergic (hypersensitive) to ingredients of the product.
- Caution if you have impaired kidney function.

-Before you start to use it

Pregnancy and lactation

Talk to your doctor before taking this medicine if you are pregnant, might become pregnant or are breastfeeding.

-Taking other medicines

Inform your doctor or pharmacist before taking this medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Antamac may interfere with each other. They may be affected by Antamac or may affect how well it works.

- It may interfere with the absorption of Tetracycline (antibiotic), Anticholinergics (medicine that block the action of acetylcholine), Barbiturates (to treat seizures), Digoxin (to treat heart problem), Warfarin (to treat blood clots) & Vitamins when taken concomitantly.

- An Interval of 3 hours is required to prevent this interference.

How to use Antamac

-How much to use

Always use Antamac exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

The usual dose:

Adults and Children above 12 years:

5-10 ml 4 times a day, after or between meals and bedtime or as directed by doctor.

Children below 12 years: 2.5ml-5ml 4 times a day, after or between meals and bedtime or as directed by doctor.

-When to use it

To be taken 4 times daily, after or between meals and bedtime. Use as directed by your doctor or pharmacist.

-How long to use it

Continue taking Antamac as your doctor recommended. If symptoms persist, consult your doctor.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much Antamac may cause:

- Hypermagnesemia (high level of magnesium in the blood).

While you are using Antamac

-Things you must do

- Tell all the doctors, dentists and pharmacists treating you that you are taking Antamac.

- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not take more than the recommended dose unless your pharmacist or doctor tells you.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give Antamac to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

- Be careful driving or operating machinery until you know how Antamac affects you.
- Do not take alcoholic beverages with this medicine.

Side effects

Like all medicines, Antamac can cause side effects, although not everybody gets them.

If side effects persist, do not take the medicine and refer to the doctor.

Tell your doctor if you notice any of the following side effects or notice any other effects not listed:

- Mild constipation/diarrhea
- Nausea
- Vomiting
- Stomach cramps
- Unusual increased in thirst

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.moh.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and disposal of Antamac

-Storage

Keep container tightly closed. Store below 30°C. Protect from light. Keep medicine out of reach of children.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

ANTAMAC SUSPENSION

Magnesium Trisilicate/Aluminium Hydroxide/Simethicone (500mg/250mg/25mg per 5ml)

Product description

-What it looks like

White Suspension with peppermint flavor

-Ingredients

-Active ingredient

Each 5ml contains

Magnesium Trisilicate (500mg),
Aluminium Hydroxide (250mg) and
Simethicone (25mg).

-Inactive ingredients

Peppermint Oil, Sodium Methyl
Paraben, Sodium Propyl Paraben,
Cetylpyridinium Chloride, Xanthan
gum, Disodium EDTA, Hydrogenated
castor oil/Kolliphor RH 40, Sorbitol
70% solution, Sucralose, Citric Acid
Anhydrous, Hydrogen Peroxide 35%,
Purified water.

-MAL number

MAL19962478X

Manufacturer and Product

Registration Holder

KCK Pharmaceutical Industries Sdn.

Bhd.

Plot 61, Bayan Lepas Industrial Park

Phase 4, Lintang Bayan Lepas 1, 11900

Bayan Lepas, Pulau Pinang, Malaysia.

Date of revision

25/11/2025