
ANTAMAC SUSPENSION

Each 5ml contains:

Aluminium Hydroxide Gel BP	250mg
Magnesium Trisilicate BP.....	500mg
Simethicone USP.....	25mg

Preservative:

Sodium Propyl Paraben.....	0.08% w/v
Sodium Methyl Paraben.....	0.14% w/v
Cetylpyridinium Chloride.....	0.1% w/v
Chloroform.....	0.25% v/v

Product Description

White suspension with peppermint flavour.

Pharmacodynamics

Small amounts of the Aluminium Hydroxide are absorbed from the intestine. Approximately 10% of the Magnesium Trisilicate is absorbed only in very minutes quantity.

Pharmacokinetics

Both Magnesium Trisilicate & Aluminium Hydroxide are slow acting antacid with a prolonged duration of action. They reach chemically to neutralize or buffer existing quantity of stomach acid but has no direct effect on its output. This action results in increased pH value of stomach content, provide a balance between constipating effect of Aluminium Hydroxide & the laxative effect of Magnesium Trisilicate. Simethicone acts by lowering the surface tension of gas bubbles, thereby causing them to coalesce and can be eliminated from the gastrointestinal tract. This action makes it a good antifatulence.

Indications

Antamac is used in the symptomatic relief of acid indigestion, dyspepsia, heartburn, gastric pain and flatulence.

Recommended Dose

Adults and children above 12years: 5-10ml 4 times a day, after or between meals and bedtime or as directed by doctor.

Route of Administration

Oral

Contraindications

No known contraindications

Warning and Precautions

It may interfere with the absorption of Tetracycline, Anticholinergics, Barbiturates, Digoxin, Warfarin & Vitamins when taken concomitantly. An interval of 3 hours is required to prevent this interference. Problems in pregnancy and lactation in human have not been established. Risk-benefit must be considered. It should be given with caution to patients with impaired renal function.

Interactions with Other Medicaments

Not know drug interaction

Pregnancy & Lactation

Pregnancy Risk Factor C

Side Effects

It may cause mild constipation or diarrhoea, nausea, vomiting, stomach cramps and unusual increased thirst.

Symptoms and Treatment of Overdose

Hypermagnesemia may be corrected by the intravenous injection of 10 to 20ml of 10% Calcium Gluconate injection. Dialysis and correction of fluid deficit may be value.

Effects on Ability to Drive and Use Machine

N/A

Storage Condition

Keep container tightly closed. Store below 30°C. Protect from light

Dosage Forms and Packaging Available

Oral liquid in 60ml, 90ml, 100ml, 120ml, 200ml, 240ml bottle.
3.6 L bottle (for export only).

Date of Revision:

26 November 2019

Registration Number:

MAL19962478X

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.
Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia